

Review

Pharmacogenomics in Lung Cancer Chemotherapy: A Review of What the Oncologist Should Know

CHIARA D'ANTONIO¹, ANNALISA MILANO¹, RICCARDO RIGHINI², CONCETTA ELISA ONESTI²,
MARIA BASSANELLI², ROSA FALCONE², IDA PARIS², SALVATORE LAURO² and PAOLO MARCHETTI²

*Departments of ¹Medical and Surgical Sciences and Translational Medicine and
²Molecular and Clinical Medicine, "Sapienza" University of Rome, Sant'Andrea Hospital, Rome, Italy*

Abstract. Lung cancer is the leading cause of cancer-related death around the world; the addition of chemotherapy to treatment of this disease has been shown to significantly increase progression-free survival and overall survival. Despite newer chemotherapies, it is important to personalize the care (treatment and dose) upon each single patient's susceptibility for controlling and reducing adverse side-effects, at best. The present review describes the current status of pharmacogenomics studies regarding germline DNA variants that may alter response and tolerability to chemotherapeutic agents used to treat lung cancer, including perspective studies.

Pharmacogenomics is an emerging branch of pharmacology that studies individual genetic variability in response to drugs, drug-induced gene expression modulation and the identification of new pharmacological targets based on the knowledge of genetic information. In fact, the human genome sequencing revealed that a single gene may have a number of differences in sequence at the nucleotide level, defined as polymorphism by combination of the Greek words poly (meaning multiple) and morph (meaning form). It is very important to analyze the correlation between patients' genotypes and phenotypes in order to define the relationship between individual polymorphisms and the alterations

This article is freely accessible online.

Correspondence to: D'Antonio Chiara, PhD, Department of Medical and Surgical Sciences and Translational Medicine, "Sapienza" University of Rome, Sant'Andrea Hospital, Grottarossa 1035, Rome, Italy. Tel: +39 0633775656, Fax: +39 0633776629, e-mail: dantonio.chiara@gmail.com

Key Words: Pharmacogenomic, lung cancer chemotherapy, polymorphism patents, cisplatin, gemcitabine, taxanes, pemetrexed, vinorelbine.

resulting by using drugs and to explain why the same therapy can have different results in different individuals. In particular, the aim of a pharmacogenomics analysis is singling-out any mutation that can affect the drug's therapeutic effects or the predisposition to undesired effect's occurrence (1). It is known that individual variation in drug response is due to factors such as age, sex, body weight, physical fitness, illnesses, level of liver and kidney functionality, diet, alcohol and tobacco use, but recent studies have reported that polymorphism of genes encoding proteins involved in the transport, metabolism and action of drugs influences in a more prominent way the outcome of chemotherapy (2-4) and toxicity. The problem of toxicity due to adverse drug reaction (ADR), despite being a major one, is often underestimated: it reportedly causes 7% of all hospitalizations in Europe, while in the USA it causes about 106,000 deaths and costs about 380 million dollars every year. It is estimated that the total cost to the community of damages caused by ADRs equals annually the cost of all pharmacological treatments (5). Chemotherapy-related ADRs seem to increase the global hospital cost by around 1,9% and the drug cost by 15% (6).

Such a variety of gene polymorphisms eventually reflects in a remarkable inter-individual variability both in chemotherapy response and toxicity, as no reliable and easily-evaluable markers are currently known to predict the drug treatments' efficacy and toxicity in neoplastic diseases. However, nowadays some peculiar polymorphisms have been identified that could cause significant side-effects and affect the tolerability and compliance to chemotherapy; in particular, in the present review we evaluated all studies demonstrating that genetic variants of polymorphisms significantly change the medical outcome in response to the administered drug in the Non Small Cell Lung Cancer (NSCLC). The patented methodologies, presented herein shed light on their identification and, consequently, mode of clinical approach. Thanks to developments in molecular

genetics, there are currently several techniques that allow the identification of known genetic alterations directly, by examining the differences in the DNA nucleotide sequence. In particular, semi-automated methods have been patented for genotyping, based on the polymerase chain reaction (PCR), permitting the identification of polymorphisms in a very simple and fast way. For instance, a 2006 patent allows analysis of the fragments by an automated capillary electrophoresis system that can separate, detect and analyse up to 16 capillaries of fluorescently labelled DNA fragments in one run (7); another patent in 2002 defined a method of sequencing DNA, based on the detection of base incorporation by the release of pyrophosphate (PPi) (8), while an older patent application, nowadays in use, describes the methodology of allelic discrimination based on chemistry (9). A more recent invention provides a method to produce a reduced representation of a genome for sequencing and DNA polymorphism detection (10).

Pharmacogenomics Analysis: Rationale

The ultimate purpose of a pharmacogenomics analysis is to carry-out a preliminary analysis of the individual patient's genetic characteristics to predict the therapeutic effect and toxicity of the employed drug, thus allowing for optimization and personalization of the chemotherapy treatment (11). Such a personalization is surely a prominent goal especially in the oncological field. For instance, as several studies have shown, the administration of the same dose of an anti-blastic drug in a patient population often results in a wide toxicity range with lethal consequences in some cases (12). Moreover, such drugs have a low therapeutic index, *i.e.* the ratio between the lowest effective and the highest tolerated doses, the latter of which is generally employed; this entails a high risk of developing adverse effects in the sub-population of genetically predisposed individuals that show an altered drug metabolism (13). Gene characterization of each patient would surely help identify the most suitable treatment protocol (14).

To this day, the dosing of chemotherapy drugs is still based on the patient's body surface correlated with the circulating blood volume and glomerular filtration rate, but not with their pharmacokinetics. Cancer patients are practically treated with a trial-and-error approach, by generally applying the treatment and operating a standard dose reduction in case of toxicity (15).

Lastly, recent studies have shown that the number of polymorphisms in various genes affects therapy efficacy (16). Drugs are involved in a series of complex metabolic pathways, in which several proteins intervene, any of which may carry mutations that can cause an alteration of its effect; the interaction among all such polymorphisms crucially conditions the treatment's final outcome.

Methods

We performed a computerized systematic literature search in electronic databases: PubMed, Embase, Cochrane Library, SCOPUS, OVID, Springer.

Studies eligible for this analysis were updated using the search terms, NSCLC, chemotherapy, patents, polymorphism, pharmacogenomics analysis and GSTP1, XRCC1, ERCC1, MRP2, MDR1, ABCB1, TTSE1, MTHFR, CYP. The inclusion criteria were as follows: (i) Only patients with advanced NSCLC were considered; (ii) All trials had to include a treatment of platinum-based agent, *i.e.* gemcitabine, vinorelbine, taxanes, pemetrexed; (iii) Only studies of outcome and toxicity related to polymorphism were discussed and not studies on the correlation risk of developing lung cancer due to single polymorphisms; (iv) Only trials reported in English were included; (v) Patents were downloaded from: <http://www.delphion.com/cgi-bin/patsearch>, www.google.com/patents, www.uspto.gov, www.freepatentsonline.com, and www.wipo.int/pctdb/en/search-simp.jsp, www.freshpatents.com.

Chemotherapeutic Agents in Lung Cancer and Polymorphic Variants Correlated with Their Metabolism and Effect

Cisplatin

The activity of platinum (pt) derivatives can be greatly affected by polymorphisms that can alter the activity or expression level of proteins intervening in the biochemical ways in which this class of antineoplastic drug is involved. Both the activity and toxicity of platinum derivatives can be affected by polymorphisms in genes that encode proteins in the following three processes.

Transmembrane Drug Transport

Most implied in the efflux mechanism of platinum derivatives is MRP2 (Multidrug resistance-associated protein 2) from the superfamily of ATP-binding cassette (ABC) proteins, whose components are integral membrane glycoproteins working as pumps. The *ABCC2* gene and the encoded protein MRP2 are normally located in hepatocytes, as well as in kidney epithelium and in intestinal enterocytes, and are, therefore, deputed to play an important role in the disposal of drugs (17). Different expression or activity of *ABCC2* gene seems to influence systemic exposure to cisplatin (18). Two polymorphisms are known: 2366C>T consisting of the substitution of a thymine for a cytosine in position 2366 in exon 18 and 4348G>A, caused by the substitution of an adenine for a guanine in position 4348 in exon 31; both variations have been observed to be possibly associated to reduced functionality of the same transporter (19). In a recent study, in advanced and metastatic lung

cancer treated with cisplatin, three functional polymorphisms of *ABCC2* (C-24T, G1249A and C3972T) were significantly associated with a better treatment response (C-24T promoter) and an increased risk of overall toxicity in particular hematological toxicity like severe thrombocytopenia (C3972T exon 28) (20). This finding was confirmed by another trial on 113 lung cancer patients treated with platinum-based chemotherapy showing the polymorphic status of MRP2 C-24T that might be a predictive marker for treatment response (21).

There is also a recent patent application disclosing the detection of polymorphism in the *ABCC2* gene that can be realized efficiently in a short time and at low cost (22).

Drug Inactivation

The inactivation process of platinum takes place through its conjugation with reduced glutathione (GSH) in a reaction catalysed by the Glutathione S-transferase (GST) enzyme, protecting the cell from damage and assisting the detoxifying process. At least three of the genes that code for GST have been shown to have functional polymorphisms and are frequently present in general populations: Glutathione S-transferase M1 (*GSTM1*), Glutathione S-transferase T1 (*GSTT1*) and Glutathione S-transferase P1 (*GSTP1*) (23). The role of GST in the detoxification of antitumor agents suggests the possible implication of GST polymorphisms to the chemotherapeutic response. In particular a GST overexpression is reported to be associated to phenomena of innate or acquired resistance to platinum derivatives; reportedly, the *GSTP1* subclass is the most involved in such a process. The main *GSTP1* polymorphism is 313 A>G, which implies the substitution in exon 5 of adenine in position 313 with a guanine. The polymorphic status of *GSTP1* 342 A>G might be the predictive marker for the treatment response of advanced NSCLC patients due to decreased detoxification of chemotherapeutic agents. In NSCLC, *GSTP1* variant genotypes (Ala/Val or Val/Val) had a significantly better survival compared to patients who had the wild type genotype (Ala/Ala; $p=0.037$) (24). *GSTP1* also seems to influence the toxicity of chemotherapy: patients carrying the homozygous mutant *GSTP1* GG genotype were at considerable risk for severe platinum-associated polyneuropathy (18% vs. 3% in wild-type vs. heterozygous mutant patients, respectively; $p=0.01$) (25); however, *GSTP1**B and *GSTP1* 105 Val haplotypes were associated, in a statistically significant way, with toxicity, and particularly with neutropenia reduction (26). Patients with deficient-type *GSTM1* were superior responders to platinum drugs than those carrying wild-type *GSTM1* ($p=0.014$) (27). The *GSTT1* $-/-$ and *GSTM1* null genotype were significantly associated with overall survival and found to be independent prognostic factors for shorter lung cancer survival (28-29). A 2004 patent application relates to a high throughput assay

for detecting the presence of clinically significant GST polymorphic alleles in a patient (30).

Mechanisms of Platinum-DNA Adducts Repair

DNA repair occurs in three important ways: by excision of the damaged base with Base-Excision Repair (BER), by excision of the nucleotide with Nucleotide-Excision Repair (NER), and by repair of the mismatches with Mismatches Repair (MMR). The DNA repair protein XRCC1 is an important component of the BER system. The NER process involves a complex of binding proteins such as Xeroderma pigmentosum complementation group C (XPC), Xeroderma pigmentosum complementation group A (XPA) and replication protein A (RPA) recruiting the Xeroderma pigmentosum complementation group D (XPD) enzyme, also known as excision repair cross-complementing group 2 (ERCC2). Also relevant are the proteins allowing for the removal of the damaged thread, such as Xeroderma pigmentosum complementation group G (XPG) and the Xeroderma pigmentosum complementation group F (XPF)/Excision repair cross-complementing group 1 (ERCC1) complex (31).

Both the BER and NER mechanisms are involved in the repair process of all Pt-DNA adducts, those produced by cisplatin as well as carboplatin or oxaliplatin. The MMR system, on the other hand, is capable of discriminating between different types of adducts, thereby providing the basis for the efficiency of oxaliplatin in cisplatin/carboplatin-resistant tumors (32).

X-Ray Repair Cross-Complementing Group 1 (XRCC1)

The XRCC1 protein is critical for repairing DNA damage induced by the platinum-based anticancer drugs, suggesting that XRCC1-mediated DNA repair capacity may markedly impact the efficacy of platinum-based therapy against NSCLC; thus, it is possible that XRCC1 could be a future predictive marker of response to treatment in advanced-stage disease patients (33).

Two principle polymorphisms have been described for the encoding gene of XRCC1: the main one consists of a substitution of an adenine for a guanine in position 28152 (28152 G>A) in exon 10, which leads to amino-acidic substitution of a glutamine for an arginine in codon 399 (Arg399Gln) (34) and Arg194Trp polymorphism.

The Arg399Gln polymorphism shows an allele frequency of 14-39% based on the population considered; in particular, in the Caucasian population the varying allele shows a 32-36% frequency (35). Such a polymorphism involves the interaction domain of XRCC1, thus altering its capability to assemble the complex needed to efficiently repair the lesion. Several studies have in fact shown that the Arg399Gln variant is associated with increased levels of DNA damage, clearly due to a decreased capability of the mutated enzyme

to repair lesions (36). A study performed on 103 patients with NSCLC treated with a platinum-based therapy has shown that polymorphism-carrying patients have a lower survival rate (37). In the BIO-FAST Trial a polymorphism in homozygous XRCC1 (Arg399Gln) showed a prognostic role in patients treated with first line based-cisplatin chemotherapy (38). In a recent meta-analysis, *XRCC1*399Gln was less favorably associated with both response rate and overall survival to chemotherapy (39).

Other polymorphisms are also involved in chemotherapy toxicity: patients carrying at least one variant of the XRCC1 Arg399Gln allele have a 2.5-fold increased risk of grade 3 or 4 gastrointestinal toxicity when treated with first-line cisplatin-based chemotherapy (40).

Yuan *et al.* found that the *XRCC1* Arg194Trp allelic variant in Asian patients was particularly associated with the response to platinum-based therapy (41).

Excision Repair Cross-Complementing Group 1 (ERCC1)

ERCC1 is a component of the NER system. Several studies have shown that this protein's expression level should apparently be correlated to the cell sensitivity to the platinum derivatives' activity. An International Adjuvant Lung Cancer Trial (IALT) understudy showed cisplatin-based adjuvant therapy to significantly prolong the survival rate in ERCC1-negative patients (42). A study by Lord *et al.*, in patients with locally-advanced or metastatic disease, showed that a low ERCC1 expression is associated to a significant improvement of the survival rate in patients treated with cisplatin and gemcitabine (43). Cobo *et al.* showed that *ERCC1* mRNA levels predict the response to cisplatin-based treatment in metastatic patients (44). In patients receiving adjuvant chemotherapy after surgical resection for lung neoplasms, the relapse rate was lower in *ERCC1*-positive subjects compared to *ERCC1*-negative, and in patients with C8092 A polymorphism. Frequent *ERCC1* polymorphism is the variation of an adenine for a cytosine in position 8092 (8092C>A); this variant shows an allele frequency of 27%. This polymorphism seems as well to have a good predictive value on a platinum-derivative-based therapy outcome. In a study by Zhou *et al.*, performed on 128 patients with advanced-stage NSCLC treated with a platinum-derivative-based therapy (cisplatin/carboplatin), individuals with wild-type C/C genotype had better survival rates (45). In another study, carriers of at least one polymorphic allele showed higher gastrointestinal toxicity with chemotherapy (46). Also in multivariate analysis, *ERCC1* expression and C8092A polymorphism were independent prognostic factors in chemotherapy-naive stage I patients (47).

One of the *ERCC1* polymorphisms is 19007 C>T, due to the transition from cytosine to thymine in position 19007, leading to alteration of a codon that encodes for the same

amino acid asparagine (Asn) but is translated at a slower speed, thus reducing the gene expression. This polymorphism has a 55-60% frequency (48) and, affecting *ERCC1* expression, can play a positive predictive role on a platinum-derivative-based therapy outcome. In a study on 109 NSCLC patients, carriers of this variant were found to have a better survival rate (49).

There are patents providing compositions and methods to analyze polymorphisms in the *ERCC1* and *XRCC1* genes for determining a patient's cancer risk and treatment response (50-51).

Methylenetetrahydrofolate Reductase (MTHFR)

Methylenetetrahydrofolate reductase (MTHFR) is an enzyme involved in the transformation of 5-10 methylenetetrahydrofolate to 5 methyltetrahydrofolate, needed as a methyl giver for the remethylation of homocysteine in methionine with the intervention of vitamin B12. Alongside severe MTHFR deficiency, another common gene polymorphism due to the substitution of a thymine for a cytosine in nucleotide 677 (C677T) causing substitution of a valine for an alanine in the final protein and a 50% reduction of MTHFR enzyme activity, has been identified. Such a variant entails high homocysteine levels in the bloodstream, especially after the administration of methionine (52).

Another mutation in the *MTHFR* gene, known as genetic variant A1298C, consists of the transition from adenine to cytosine in position 1298 leading to the substitution of a glutamate with an alanine. This genetic variant is associated with high homocysteine levels and low folate levels in plasma when combined with the C677T mutation (53). MTHFR has been related to response at platinum-based chemotherapy in NSCLC. A recent meta-analysis confirmed that MTHFR 677TT homozygote carriers had a better response to platinum-based chemotherapy. Patients in treatment with homozygous mutations for *MTHFR* C677T had a significantly increased PFS (progression free survival) compared to patients with wild-type or heterozygous mutations (54). A single nucleotide polymorphism of *MTHFR* has also been associated with an increased risk of toxicity, like irradiation pneumonia developing in lung cancer patients treated with radiation therapy to the chest wall (55).

Finally, there is finally a patent disclosing a primer set for specifically amplifying a target region in the *MTHFR* gene by a nucleic acid amplification method (56).

Gemcitabine

Although many reports about single nucleotide polymorphisms (SNPs) in the family of ABC transporters have been published, the impact of polymorphisms on pharmacokinetics and pharmacodynamics of gemcitabine remains to be defined (57). More studies report multidrug resistance 1 (MDR1) overexpression to be associated with

sensitivity to gemcitabine (58). *MDR1* plays a major role in drug resistance by impairing the intracellular retention of multiple anticancer drugs. The most studied SNP is the C3435T located in exon 26. This polymorphism has been found to have a role in the function of the permeability glycoprotein (P-gp) responsible for substance elimination by the cell membrane by hydrolysis of ATP; multiple studies showed that this polymorphism was significantly correlated with drug response (58). The overall response to treatment of homozygotes for the wild-type (C/C) allele in *MDR1* C3435T was significantly better than in heterozygotes (C/T) (59) and mutant homozygotes (T/T) (95% confidence interval (CI): 1.44-3.68; $I=0.0005$) (60). The *MDR1* 2677 polymorphism seems to be correlated with an increase in grade 3 toxicities in the mutated group (39%) and an increase in both progression-free and global survival in patients that had had gemcitabine as an adjuvant for other cancers (61).

Of note, there is a patent for determining haplotypes or diplotypes of the 5' regulatory region of *MDR1* (multidrug resistance 1) gene (62). Another recent patent consists of a single nucleotide polymorphism (SNP) detection kit for G2677T/A of *MDR1* (multidrug resistance gene 1) (63).

Taxanes

The overexpression of ABC-transporter genes such as *MDR1*, is a factor of non-response to taxane-based chemotherapy due to increased toxicity originating from a longer exposure to the drug, due, in turn, to a reduced elimination rate (64). Overexpression of P-gp1/*ABCB1* is in fact associated with poor prognosis in several tumor types. Moreover, polymorphisms in cytochromes *P450* (3 A4, 3 A5, 2 C8), that are implied in the hepatic metabolism of taxanes, can determine an increase in drug concentration due to lack of elimination and thus an unacceptable toxicity level (65). Both genes *CYP3A4* and *CYP3A5* are polymorphic and several allele variants have been described. Only two allele variants that are in linkage disequilibrium, namely *CYP3A4*1B* and *CYP3A5*3*, are common to several ethnicities and have functional relevance (66). As far as the *CYP3A4*1B* is concerned, it seems to modify the ability to metabolize some *CYP3A* substrates. *CYP3A5*3* instead seems to be the most common *CYP3A5* allele and is associated with decreased enzyme activity. In a study published in the Journal of Clinical Oncology (JCO), gaps in allelic distribution for genes involved in paclitaxel disposition or DNA repair between Japanese and US patients were observed. In this exploratory analysis, genotype-related associations with patient outcomes (progression free survival) were observed for *CYP3A4*1B* ($p=0.04$), however, this association should be interpreted in the context that only African-American patients harbored this allele (67). A 2010 patent describes a method for predicting a subject's response

to at least one *CYP3A4*-metabolized compound; with the method comprising the detection of the allelic status of one or more polymorphisms in a nucleic acid sample of the subject (68).

Pemetrexed

Pemetrexed (Alimta) is a multi-target agent converted to a series of polyglutamate-derived metabolites by folylpolyglutamate synthetase. They inhibit three folate-dependent enzymes: thymidylate synthase (TS), dihydrofolate reductase (DHFR) and glycinamide ribonucleotide formyltransferase (GARF1) (69). Several studies are focused on the effects of pemetrexed on thymidylate synthase. TS is an important target for pemetrexed-based chemotherapy. Its overexpression is correlated to TS-target chemotherapy resistance. Polymorphic tandem repeats located in the TS enhancer region (TSER) have been shown to influence the expression of TS. Three-fold tandem repeats (TSER*3) give larger *in vitro* TS expression than two-fold tandem repeats. In a Korean study, survival was significantly longer with pemetrexed in patients with high TS expression genotypes (second Nief's classification) (70) 3RGCC/3RGCC or 3RGGC/3RGGC compared to the other groups (PFS: 5.2 months vs. 3.7 months, $p=0.03$; OS: 31.8 months vs. 18.5 months, $p=0.001$) (71). In advanced non-small cell lung cancer patients treated with pemetrexed-based chemotherapy with the TS 2R/2R, 2R/3C or 3C/3C genotypes, median PFS times and response rate were significantly longer than those of patients with the 2R/3G, 3C/3G or 3G/3G genotypes ($I=0.036$ and $I=0.044$, respectively) (72). Prospective data are needed to confirm the impact of TSER on the outcome after TS-target therapy (73). There is a recent patent directed to a method of predicting a response to a chemotherapeutic regimen based on loss of heterozygosity at the thymidylate synthase locus in cancer tissue (74).

A 2009 study published by JCO on patients treated with Alimta in a second-line treatment for lung cancer has investigated polymorphisms in *MTHFR*. Homozygous mutant patients with *MTHFR* C677T have a significant increase in PFS compared to wild-type homozygous mutant patients, while homozygous mutant patients with *MTHFR* A1298C have shorter PFS as a trend ($p=0.06$) (75). In an Italian study, a multivariate analysis confirmed the independent prognostic significance of *MTHFR*-C677T both in risk of disease progression (CC-CT genotypes hazard ratio [HR] 1.94, 95%CI: 1.15-3.28; $p=0.012$) and death (HR 2.00, 95% CI: 1.12-3.54; $p=0.018$) (76).

Vinorelbine

Vinorelbine is a semi-synthetic vinca alkaloid, mainly metabolized by *CYP3A4* and *CYP3A5* (77). *CYP3A5* is a highly polymorphic gene: the widespread allele *CYP3A5*3* causes a reduced expression, hence a lower drug

Table I. Portrayal of the most important polymorphisms identified, their functional effect, the anticancer drugs involved and expected event(s).

Gene polymorphism	Functional effect	Anticancer drugs involved	Observed event
MRP2 C2366T (16)	T allele is associated with decreased transporter activity	Cisplatin	Polymorphism associated with better treatment response
MRP2 C4348A (16)	A allele is associated with decreased transporter activity	Cisplatin	Polymorphism associated with better treatment response
MRP2 C-24T (17)	T allele is associated with decreased transporter activity	Cisplatin	Polymorphism associated with better treatment response
MRP2 C3972T (17)	T allele is associated with decreased transporter activity	Cisplatin	Polymorphism associated with better treatment response and increased toxicity
MRP2 G1249A (17)	A allele is associated with decreased transporter activity	Cisplatin	Polymorphism associated with better treatment response
GSTP1 A313G (20)	G allele is associated with decreased enzyme activity	Cisplatin	Polymorphism associated with better survival and increased toxicity
GSTP1 A342G (19)	G allele is associated with decreased enzyme activity	Cisplatin	Polymorphism associated with better survival and increased toxicity
GSTM1 gene deletion (23)	Loss of function	Cisplatin	Polymorphism associated with better response to platinum drugs
GSTT1 gene deletion (24)	Loss of function	Cisplatin	Polymorphism associated with shorter survival
XRCC1 G28152A (31-37)	A allele is associated with decreased DNA repair activity	Cisplatin	Polymorphism associated with lower OS and response rate
ERCC1 C8092A (41-44)	A allele is associated with decreased DNA repair activity	Cisplatin	Polymorphism associated with better survival and increased toxicity
ERCC1 C19007T (45-46)	C allele is associated with decreased DNA repair activity	Cisplatin	Polymorphism associated with better survival
MTHFR C677T (48-52,70-74)	T allele is associated with decreased enzyme activity	Cisplatin, pemetrexed	Polymorphism associated with better response to platinum; Homozygous mutant shows increased PFS
MTHFR A1298C (49, 70)	C allele is associated with decreased enzyme activity	Cisplatin, pemetrexed	Homozygous mutant has shorter PFS
MDR1 C3435T (56-57,77)	T allele is associated with decreased transporter activity	Gemcitabine, vinorelbine	Polymorphism associated with worse response to gemcitabine; Lower risk of progression in patients treated with Vinorelbine
MDR1 G2677T (80)	T allele is associated with decreased transporter activity	Gemcitabine, vinorelbine	Polymorphism associated with increased toxicities and better OS and PFS in patients treated with Gemcitabine; Better response to Vinorelbine
CYP3A4*1B (89)	Polymorphism associated with decreased enzyme activity	Paclitaxel, vinorelbine	Polymorphism associated with response
CYP3A5*3 (88, 73)	Polymorphism associated with decreased enzyme activity	Paclitaxel, vinorelbine	Polymorphism associated with response
TTSER 28 bp VNTR (2R/3R) (68)	3R allele is associated with increased TS expression (increased enzyme activity)	Pemetrexed	Survival observed was significantly longer in patients with high expression genotype

metabolism in the liver (78). In a study by Pan *et al.* in 59 NSCLC patients, the CYP3A5*3 polymorphism seemed to be associated to responses to vinorelbine treatment. No significant difference in toxicity and survival was observed according to SNP genotype in lung cancer patients (79). Vinorelbine is also a substrate for P-gp membrane transporters (MDR1, ABCB1). *MDR1* polymorphisms can cause alterations in the drug's absorption and elimination

(80). *MDR1* 3435CC polymorphisms appear to be related to lower risk of progression in advanced non small cell cancer treated with vinorelbine (81). In a 2008 study published in *Respiration*, *ABCB1* C3435T polymorphism was associated to a better response to vinorelbine-based chemotherapy for lower P-gp expression levels (82). This study was in agreement with some reports but in disagreement with others, thus further studies should be

carried on with large patient cohorts to confirm these results (83-84). There is a patent disclosing the detection of this gene polymorphism having an influence on pharmacokinetics in a DNA sample from a subject and predicting the influence of the gene polymorphism on the kinetics of a drug in the subject (85).

Conclusion

Pharmacogenomics provide a polygenic, global approach in the genome study, by simultaneously analysing multiple gene polymorphisms or multiple mutations within a single gene. The pharmacogenomics analysis, aimed at inquiring the effect of single polymorphisms on the treatment's outcome is still fundamental in situations in which the alteration of even a single gene can play a crucial role, such as the predisposition to developing toxic effects.

Integrating results from pharmacogenetics and pharmacogenomics studies surely is a valuable strategy to be developed in the coming years in order to finally achieve individualization of the anti-blastic therapy and thus obtain the highest efficacy and lowest toxicity from each treatment. Polymorphic variants of enzymes involved in the metabolism of chemotherapeutic agents used in lung cancer should be investigated (Table I).

Current and Future Developments

Randomized clinical trials are required in order to define not only the correlation with toxicity, but also with the outcome. In fact, the effectiveness of therapy may be influenced by genetic characteristics of the tumor cells because of their high genetic instability often developing additional somatic genetic alterations that lead to genotype differences that are not found in non-neoplastic germ cells. The pharmacogenomics studies, on the other hand, are based mainly on an analysis of the germ genetic characteristics obtained from peripheral blood samples. Thus, the purpose of these studies is to define a personalized therapy, which gives the maximum clinical benefit with minimal occurrence of side effects. Today, such an approach seems as a distant goal because it is necessary to consider various factors that may influence the chemotherapy response, such as mode of drug administration, administration of other drugs/chemotherapeutic agents, demographic and clinical characteristics of patients, ethnicity issues, influence of environmental factors (alcohol, smoking, diet), all of which interfere with the standardization of the data obtained from this type of analysis.

Conflicts of Interest

The Authors declare that they have no conflicts of interest.

References

- 1 Brunton LL, Parker K L, Blumenthal D and Buxton I: The Goodman & Gilman's manual of pharmacology and therapeutics. 11th Edition. The McGraw-Hill Companies: New York 2008.
- 2 Nagasubramanian R, Innocenti F and Ratain MJ: Pharmacogenetics in cancer treatment. *Annu Rev Med* 54: 437-452, 2003.
- 3 Relling MV and Dervieux T: Pharmacogenetics and cancer therapy. *Nat Rev Cancer* 1(2): 99-108, 2001.
- 4 Innocenti F and Ratain MJ: Update on pharmacogenetics in cancer chemotherapy. *Eur J Cancer* 38(5): 639-644, 2002.
- 5 Lazarou J, Pomeranz BH and Corey PN: Incidence of adverse drug reactions in hospitalized patients: a meta-analysis of prospective studies. *JAMA* 279(15): 1200-1205, 1998.
- 6 Johnson JA and Bootman JL: Drug-related morbidity and mortality. A cost-of-illness model. *Arch Intern Med* 155(18): 1949-1956, 1995.
- 7 Heiner C, Nordman ES and Woudenberg TM: Compositions and methods for clonal amplification and analysis of polynucleotides. WO 2006099579 A2, 2006.
- 8 Nyren P, Ronaghi M and Tallisjoe A: Method of sequencing DNA. WO 2002020836 A2, 2002.
- 9 Gelfand DH, Holland PM, Saiki RK and Watson RM: Reaction mixtures for detection of target nucleic acids. US 5804375 A, 1998.
- 10 Kirst M, Ribeiro De Resende MF, JR, Gomide Neves L, Dervinis C and Balmant KM: Method for genome complexity reduction and polymorphism detection. US 20130035238 A1, 2013.
- 11 Giovannetti E, Toffalorio F, De Pas T and Peters GJ: Pharmacogenetics of conventional chemotherapy in non small cell lung cancer: a changing landscape? *Pharmacogenomics* 13(9): 1073-1086, 2012.
- 12 Ferté C, Gomez Roca C, Loriot Y, Bahleda R, Moldovan C, Cohen A, Izzedine H, Soria JC and Ederhy S: Reversible cardiogenic shock following 5-fluorouracil infusion. *Invest New Drugs* 28(4): 531-533, 2010.
- 13 Yamayoshi Y, Iida E and Tanigawara Y: Cancer pharmacogenomics: international trends. *Int J Clin Oncol* 10(1): 5-13, 2005.
- 14 Marck PC, Gandara DR and Lara PN: Efficacy and toxicity differences in lung cancer populations in the era of clinical trials globalization: the "common arm" approach. *Expert Rev Anticancer Ther* 12(12): 1591-1596, 2012.
- 15 Evans WE and McLeod HL: Pharmacogenomics-drug disposition, drug targets, and side effects. *N Engl J Med* 348(6): 538-549, 2003.
- 16 Marsh S and McLeod HL: Cancer pharmacogenetics. *Br J Cancer* 90(1): 8-11, 2004.
- 17 Lockhart AC, Tirona RG and Kim RB: Pharmacogenetics of ATP-binding cassette transporters in cancer and chemotherapy. *Mol Cancer Ther* 2(7): 685-698, 2003.
- 18 Cascorbi I and Haenisch S: Pharmacogenetics of ATP-binding cassette transporters and clinical implications. *Methods Mol Biol* 596: 95-121, 2010.
- 19 Hirouchi M, Suzuki H, Itoda M, Ozawa S, Sawada J, Ieiri I, Ohtsubo K and Sugiyama Y: Characterization of the cellular localization, expression level, and function of SNP variants of MRP2/ABCC2. *Pharm Res* 21(5): 742-748, 2004.

- 20 Han B, Gao G, Wu W, Gao Z, Zhao X, Li L, Qiao R, Chen H, Wei Q, Wu J and Lu D: Association of ABCC2 polymorphisms with platinum-based chemotherapy response and severe toxicity in non small cell lung cancer patients. *Lung Cancer* 72(2): 238-243, 2011.
- 21 Sun N, Sun X, Chen B, Cheng H, Feng J, Cheng L, Fenq J, Cheng L and Lu Z: MRP2 and GSTP1 polymorphisms and chemotherapy response in advanced non-small cell lung cancer. *Cancer Chemother Pharmacol* 65(3): 437-446, 2010.
- 22 Iguchi A: Probe, polymorphism detection method, method of evaluating drug efficacy or tolerance, and reagent kit. US 20120282608 A1, 2012.
- 23 Joerger M, Burgers SA, Baas P, Smit EF, Haitjema TJ, Bard MP, Doodeman VD, Smits PH, Vincent A, Huitema AD, Beijnen JH and Schellens JH: Germline polymorphisms in patients with advanced non-small cell lung cancer receiving first-line platinum-gemcitabine chemotherapy: a prospective clinical study. *Cancer* 118(9): 2466-2475, 2012.
- 24 Lu C, Spitz MR, Zhao H, Dong Q, Truong M, Chang JY, Blumenschein GR Jr, Hong WK and Wu X: Association between glutathione S-transferase polymorphisms and survival in patients with advanced non-small cell lung carcinoma. *Cancer* 106(2): 441-447, 2006.
- 25 Booton R, Ward T, Heighway J, Aschcroft L, Morris J and Thatcher N: Glutathione-S-transferase P1 isoenzyme polymorphisms, platinum-based chemotherapy, and non small cell lung cancer. *J Thorac Oncol* 1(7): 679-683, 2006.
- 26 Li W, Yue W, Zhang L, Zhao X, Ma L, Yang X, Zhang C, Wang Y and Gu M: Polymorphisms in GSTM1, CYP1A1, CYP2E1, and CYP2D6 are associated with susceptibility and chemotherapy response in non-small-cell lung cancer patients. *Lung* 190(1): 91-98, 2012.
- 27 Sreeja L, Syamala V, Hariharan S, Syamala VS, Raveendran PB, Sivanandan CD, Madhavan J and Ankathil R: Glutathione S-transferase M1, T1 and P1 polymorphisms: susceptibility and outcome in lung cancer patients. *J Exp Ther Oncol* 7(1): 73-85, 2008.
- 28 Gonlugur U, Pinarbasi H, Gonlugur TE and Silig Y: The association between polymorphisms in glutathione S-transferase (GSTM1 and GSTT1) and lung cancer outcome. *Cancer Invest* 24(5): 497-501, 2006.
- 29 Sweeney C, Nazar-Stewart V, Stapleton PL, Eaton DL and Vaughan TL: Glutathione S-transferase M1, T1, and P1 polymorphisms and survival among lung cancer patients. *Cancer Epidemiol Biomarkers Prev* 12(6): 527-533, 2003.
- 30 Ali-Osman F, Ballard L, Keller C and Lemmons R: High throughput detection of glutathione s-transferase polymorphic alleles. WO 2004036179 A2, 2004.
- 31 Marsin S, Vidal AE, Sossou M, Menissier-de Murcia J, Le Page F, Boiteux S, de Murcia G and Radicella JP: Role of XRCC1 in the coordination and stimulation of oxidative DNA damage repair initiated by the DNA glycosylase hOGG1. *J Biol Chem* 278(45): 44068-44074, 2003.
- 32 Raymond E, Faivre S, Chaney S, Woinarowski J and Cvitkovic E: Cellular and molecular pharmacology of oxaliplatin. *Mol Cancer Ther* 1(3): 227-235, 2002.
- 33 Sun X, Li F, Sun N, Shukui Q, Baoan C, Jifeng F, Lu C, Zuhong L, Hongyan C, YuanDong C, Jiazhong J and Yingfenq Z: Polymorphisms in XRCC1 and XPG and response to platinum-based chemotherapy in advanced non-small cell lung cancer patients. *Lung Cancer* 65(2): 230-236, 2009.
- 34 Duell EJ, Holly EA, Bracci PM, Whiencke JK and Kelsey KT: A population-based study of the Arg399Gln polymorphism in Xray repair cross- complementing group 1 (XRCC1) and risk of pancreatic adenocarcinoma. *Cancer Res* 62(16): 4630-4636, 2002.
- 35 Goode EL, Ulrich CM and Potter JD: Polymorphisms in DNA repair genes and associations with cancer risk. *Cancer Epidemiol Biomarkers Prev* 11(12): 1513-1530, 2002.
- 36 Wang Y, Spitz MR, Zhu Y, Dong Q, Shete S and Wu X: From genotype to phenotype: correlating XRCC1 polymorphisms with mutagen sensitivity. *DNA Repair (Amst)* 2(8): 901-908, 2003.
- 37 Gurubhagavatula S, Liu G, Park S, Zhou W, Su L, Wain JC, Lynch TJ, Neuberger DS and Christiani DC: XPD and XRCC1 genetic polymorphisms are prognostic factors in advanced non-small-cell lung cancer patients treated with platinum chemotherapy. *J Clin Oncol* 22(13): 2594-2601, 2004.
- 38 Tiseo M, Bordi P, Bortesi B, Boni L, Boni C, Baldini E, Grossi F, Recchia F, Zanelli F, Fontanini G, Naldi N, Campanini N, Azzoni C, Bordi C and Ardizzoni A: Bio-FAST trial Group: ERCC1/BRCA1 expression and gene polymorphisms as prognostic and predictive factors in advanced NSCLC treated with or without cisplatin. *Br J Cancer* 108(8): 1695-1703, 2013.
- 39 Wu J, Liu J, Zhou Y, Ying J, Zou H, Guo S, Wang L, Zhao N, Hu J, Lu D, Jin L, Li Q and Wang JC: Predictive value of XRCC1 gene polymorphisms on platinum-based chemotherapy in advanced non-small cell lung cancer patients: a systematic review and meta-analysis. *Clin Cancer Res* 18(14): 3972-3981, 2012.
- 40 Wang Z, Xu B, Lin D, Tan W, Leaw S, Hong X and Hu X: XRCC1 polymorphisms and severe toxicity in lung cancer patients treated with cisplatin-based chemotherapy in Chinese population. *Lung Cancer* 62(1): 99-104, 2008.
- 41 Yuan P, Miao XP, Zhang XM, Wang ZH, Tan W, Sun Y, Zhang XR, Xu BH and Lin DX: XRCC1 and XPD genetic polymorphisms predict clinical responses to platinum-based chemotherapy in advanced non-small cell lung cancer. *Zhonghua Zhong Liu Za Zhi* 28(3): 196-199, 2006.
- 42 Olausson KA, Dunant A, Fouret P, Brambilla E, André F, Haddad V, Taranchon E, Filipits M, Pirker R, Popper HH, Stahel R, Sabatier L, Pignon JP, Tursz T, Le Chevalier T and Soria JC; IALT Bio Investigators: DNA repair by ERCC1 in non-small-cell lung cancer and cisplatin-based adjuvant chemotherapy. *N Engl J Med* 355(10): 983-991, 2006.
- 43 Lord RV, Brabender J, Gandara D, Alberola V, Camps C, Domine M, Cardenal F, Sanchez JM, Gumerlock PH, Tarón M, Sánchez JJ, Danenberg KD, Danenberg PV and Rosell R: Low ERCC1 expression correlates with prolonged survival after cisplatin plus gemcitabine chemotherapy in non-small cell lung cancer. *Clin Cancer Res* 8(7): 2286-2291, 2002.
- 44 Cobo M, Isla D, Massuti B, Montes A, Sanchez JM, Provencio M, Vinolas N, Paz-Ares L, Lopez-Vivanco G, Munoz MA, Felip E, Alberola V, Camps C, Domine M, Sanchez JJ, Sanchez-Ronco M, Danenberg K, Taron M, Gandara D and Prosell R: Customizing cisplatin based on quantitative excision repair cross complementing I mRNA expression: a phase III trial in non-small-cell lung cancer. *J Clin Oncol* 25(19): 2747-2754, 2007.
- 45 Zhou W, Gurubhagavatula S, Liu G, Park S, Neuberger DS, Wain JC, Lynch TJ, Su L and Christiani DC: Excision repair cross-complementation group 1 polymorphism predicts overall survival in advanced non-small cell lung cancer patients treated with platinum-based chemotherapy. *Clin Cancer Res* 10(15): 4939-4943, 2004.

- 46 Suk R, Gurubhagavatula S, Park S, Zhou W, Su L, Lynch TJ, Wain JC, Neuberger D, Liu G and Christiani DC: Polymorphisms in ERCC1 and grade 3 or 4 toxicity in nonsmallnon small cell lung cancer patients. *Clin Cancer Res* 11(4): 1534-1538, 2005.
- 47 Tseden-Ish M, Choi YD, Cho HJ, Ban HJ, Oh IJ, Kim KS, Song SY, Na KJ, Ahn SJ, Choi S and Kim YC: Disease-free survival of patients after surgical resection of non-small cell lung carcinoma and correlation with excision repair cross-complementation group 1 expression and genotype. *Respirology* 17(1): 127-133, 2012.
- 48 Viguier J, Boige V, Miquel C, Pocard M, Girandeau B, Sabourin JC, Ducreux M, Sarasin A and Praz F: ERCC1 codon 118 polymorphism is a predictive factor for the tumor response to oxaliplatin/5-fluorouracil combination chemotherapy in patients with advanced colorectal cancer. *Clin Cancer Res* 11(17): 6212-6217, 2005.
- 49 Ryu JS, Hong YC, Han HS, Lee JE, Kim S, Park YM, Kim YC and Hwang TS: Association between polymorphisms of ERCC1 and XPD and survival in non-small-cell lung cancer patients treated with cisplatin combination chemotherapy. *Lung Cancer* 44(3): 311-316, 2004.
- 50 Lenz HJ, Stoehlmacher J and Park (Jong-Han) D: Polymorphisms in the ERCC1 gene for predicting treatment outcome. *US 7662553 B2*, 2010.
- 51 Lenz HJ, Stoehlmacher J and Park (Jong-Han) D: Polymorphisms for predicting disease and treatment outcome. *EP 2385137 A1*, 2011.
- 52 Zhao M, Li X, Xing C and Zhou B: Association of methylenetetrahydrofolate reductase C677T and A1298C polymorphisms with colorectal cancer risk: a meta-analysis. *Biomed Rep* 1(5): 781-791, 2013.
- 53 Zhu N, Gong Y, He J, Xia J and Chen X: Influence of Methylenetetrahydrofolate Reductase C677T Polymorphism on the Risk of Lung Cancer and the Clinical Response to Platinum-Based Chemotherapy for Advanced Non-Small Cell Lung Cancer: An Updated Meta-Analysis. *Yonsei Med J* 54(6): 1384-1393, 2013.
- 54 Smit EF, Burgers SA, Biesma B, Smit HJ, Eppinga P, Dingemans AM, Joerger M, Schellens JH, Vincent A, van Zandwijk N and Groen HJ: Randomized phase II and pharmacogenetic study of pemetrexed compared with pemetrexed plus carboplatin in pretreated patients with advanced non-small-cell lung cancer *J Clin Oncol* 27(12): 2038-2045, 2009.
- 55 Mak RH, Alexander BM, Asomaning K, Heist RS, Liu CY, Su L, Zhai R, Ancukiewicz M, Napolitano B, Niemirko A, Willers H, Choi NC and Christiani DC: A single nucleotide polymorphism in the MTHFR (methylene tetrahydrofolate reductase) gene is associated with risk of radiation pneumonitis in lung cancer patients treated with thoracic radiation therapy. *Cancer* 118(14): 3654-3665, 2012.
- 56 Hirai M and Kozuka M: Primer set for amplification of mthfr gene, mthfr gene amplification reagent comprising same, and use of same. *WO 2011062258 A1*, 2011.
- 57 Lepper ER, Nooter K, Verweij J, Acharya RM, Figg WD and Sparreboom A: Mechanisms of resistance to anticancer drug: the role of the polymorphic ABC transporters ABCB1 and ABCG2. *Pharmacogenomics* 6(2): 115-138, 2005.
- 58 Bergman AM, Pinedo HM, Talianidis I, Veerman G, Loves WJ, Van der Wilt CL and Peters GJ: Increased sensitivity to Gemcitabine of P-glycoprotein and multidrug resistance-associated protein-overexpressing human cancer cell lines. *Br J Cancer* 88(12): 1963-1970, 2003.
- 59 Takara K, Sakaeda T and Okumura K: An update on overcoming MDR1-mediated multidrug resistance in cancer chemotherapy. *Curr Pharm Des* 12(3): 273-286, 2006.
- 60 Wei HB, Hu J, Shang LH, Zhang YY, Lu FF, Wei M and Yu Y: A meta-analytic review of ERCC1/MDR1 polymorphism and chemosensitivity to platinum in patients with advanced non-small cell lung cancer. *Chin Med J (Engl)* 125(16): 2902-2907, 2012.
- 61 Kasuya K, Tsuchida A, Nagakawa Y, Suzuki Y, Suzuki M, Aoki T, Abe Y, Shimazu M and Sofuni A: Prediction of a side effect and efficacy of adjuvant chemotherapy with gemcitabine for post operative patient of pancreatic cancer by genetic polymorphism analysis. *Hepatogastroenterology* 59(117): 1609-1613, 2012.
- 62 Kuwano M and Wada M: SNPs in 5' regulatory region of MDR1 gene. *WO2005028645A1*, 2006.
- 63 Single nucleotide polymorphism detection kit for G2677T / A of MDR1. *CN102816845 A*, 2012.
- 64 Ross DD and Nakanishi T: Impact of breast cancer resistance protein on cancer treatment outcomes. *Methods Mol Biol* 596: 251-290, 2010.
- 65 Zhang YY, Liu Y, Zhang JW, Ge GB, Wang LM, Sun J and Yang L: Characterization of human cytochrome P450 isoforms involved in the metabolism of 7-epipaclitaxel. *Xenobiotica* 39(4): 283-292, 2009.
- 66 Xie HG, Wood AJ, Kim RB, Stein CM and Wilkinson GR: Genetic variability in CYP3A5 and its possible consequences. *Pharmacogenomics* 5(3): 243-272, 2004.
- 67 Gandara DR, Kawaguchi T, Crowley J, Moon J, Furuse K, Kawahara M, Teramukai S, Ohe Y, Kubota K, Williamson SK, Gautschi O, Lenz HJ, McLeod HL, Lara PN Jr., Coltman CA Jr., Fukuoka M, Saijo N, Fukushima M and Mach PC: Japanese-US common-arm analysis of paclitaxel plus carboplatin in advanced non-small-cell lung cancer: a model for assessing population-related pharmacogenomics *J Clin Oncol* 27(21): 3540-3546, 2009.
- 68 Sadee W and Wang D: Polymorphism in CYP3A4 gene affecting drug metabolizing and uses thereof. *WO 2010111600 A1*, 2010.
- 69 Chattopadhyay S, Moran RG and Goldman ID: Pemetrexed: biochemical and cellular pharmacology, mechanisms and clinical applications. *Mol Cancer Ther* 6(2): 404-417, 2007.
- 70 Nief N, Le Morvan V and Robert J: Involvement of gene polymorphisms of thymidylate synthase in gene expression, protein activity and anticancer drug cytotoxicity using the NCI-60 panel. *Eur J Cancer* 43(5): 955-962, 2007.
- 71 Jung M, Lee CH, Park HS, Lee JH, Kang YA, Kim SK, Chang J, Kim DJ, Rha SY, Kim JH and Cho BC: Pharmacogenomic assessment of outcomes of pemetrexed-treated patients with adenocarcinoma of the lung. *Yonsei Med J* 54(4): 854-864, 2013.
- 72 Li WJ, Jiang H, Fang XJ, Ye HL, Liu MH, Liu YW, Chen Q, Zhang L, Zhang JY, Yuan CL and Zhang QY: Polymorphisms in thymidylate synthase and reduced folate carrier (SLC19A1) genes predict survival outcome in advanced non-small cell lung cancer patients treated with pemetrexed-based chemotherapy. *Oncol Lett* 5(4): 1165-1170, 2013.
- 73 Marsh S and McLeod HL: Thymidylate synthase pharmacogenetics in colorectal cancer. *Clin Colorectal Cancer* 1(3): 179-181, 2001.
- 74 Danenberg KD: Methods of determining a chemotherapeutic regimen based on loss of heterozygosity at the thymidylate synthase locus. *CA 2517384 C*, 2013.

- 75 Smit EF, Burgers SA, Biesma B, Smit HJ, Eppinga P, Dingemans AM, Joerger M, Schellens JH, Vincent A, van Zandwijk N and Groen HJ: Randomized phase II and pharmacogenetic study of pemetrexed compared with pemetrexed plus carboplatin in pretreated patients with advanced Non Small cell lung cancer. *JCO* 27: 2038-2045, 2009.
- 76 Tiseo M, Giovannetti E, Tibaldi C, Camerini A, Di Costanzo F, Barbieri F, Burgers JA, Vincent A, Peters GJ and Ardizzoni A: Pharmacogenetic study of patients with advanced non-small cell lung cancer (NSCLC) treated with second-line pemetrexed or pemetrexed-carboplatin. *Lung Cancer* 78(1): 92-99, 2012.
- 77 Kajita J, Kuwabara T, Kobayashi H and Kobayashi S: CYP3A4 is mainly responsible for the metabolism of a new vinca alkaloid, vinorelbine, in human liver microsomes. *Durg Metab Dispos* 28(9): 1121-1127, 2000.
- 78 Chou FC, Tzeng SJ and Huang JD: Genetic polymorphism of cytochrome P450 3A5 in Chinese. *Drug Metab Dispos* 29(9): 1205-1209, 2001.
- 79 Pan JH, Han JX, Wu JM, Sheng LJ and Huang HN: CYP450 polymorphisms predict clinical outcomes to vinorelbine-based chemotherapy in patients with Non-small cell lung cancer. *Acta Oncol* 46(3): 361-366, 2007.
- 80 Woodahl EL and Ho RJ: The role of MDR1 genetic polymorphisms in interindividual variability in P-glycoprotein expression and function. *Curr Drug Metab* 5(1): 11-19, 2004.
- 81 Viñolas N, Provencio M, Reguart N, Cardenal F, Alberola V, Sánchez-Torres JM, Barón FJ, Cobo M, Maestu I, Moreno I, Mesia C, Izquierdo A, Felip E, López-Brea M, Marquez A, Sánchez-Ronco M, Tarón M, Santarpia MC and Rosell R; Spanish Lung Cancer Group : Single nucleotide polymorphisms in MDR1 gen correlates with outcome in advanced non-small-cell lung cancer patients treated with cisplatin plus vinorelbine. *Lung Cancer* 71(2): 191-198, 2011.
- 82 Pan JH, Han JX, Wu JM, Sheng LJ, Huang HN and Yu QZ: MDR1 Single Nucleotide Polymorphisms Predict Response to Vinorelbine-Based Chemotherapy in Patients with Non Small Cell Lung Cancer. *Respiration* 75(4): 380-385, 2008.
- 83 Illmer T, Schuler US, Thiede C, Schwarz UI, Kim RB, Gotthard S, Freund D, Schäkel U, Ehninger G and Schaich M: MDR1 gene polymorphisms affect therapy outcome in acute myeloid leukemia patients. *Cancer Res* 62(17): 4955-4962, 2002.
- 84 Siegsmond M, Brinkmann U, Schaffeler E, Weirich G, Schwab M, Eichelbaum M, Fritz P, Burk O, Decker J, Alken P, Rothenpieler U, Kerb R, Hoffmeyer S and Brauch H: Association of the glycoprotein transporter MDR1C3435T polymorphism with the susceptibility to renal epithelial tumors. *J Am Soc Nephrol* 13(7): 1847-1854, 2002.
- 85 Ishikawa T, Sakurai A and Tamura A: Method for detection of gene polymorphism and method for screening of substance. WO 2007026940 A1, 2007.

Received May 16, 2014

Revised July 3, 2014

Accepted July 7, 2014