

Role of Pur α in the Modulation of Homologous Recombination-directed DNA Repair by HIV-1 Tat

HUICHEN WANG¹, MARTYN K. WHITE¹, RAFAL KAMINSKI¹, NUNE DARBINIAN¹, SHOHREH AMINI^{1,2}, EDWARD M. JOHNSON³, KAMEL KHALILI¹ and JAY RAPPAPORT¹

¹*Department of Neuroscience, Center for Neurovirology, Temple University School of Medicine, Philadelphia, PA 19122;*

²*Department of Biology, College of Science and Technology, Temple University, Philadelphia, PA 19122;*

³*Department of Microbiology and Molecular Cell Biology, Eastern Virginia Medical School, Norfolk, VA 23501, U.S.A.*

Abstract. *Background: The nucleic acid-binding protein Pura is involved at stalled DNA replication forks, in double-strand break (DSB) DNA repair and the cellular response to DNA replication stress. Pura interacts with HIV-1 Tat, which regulates homologous recombination-directed DNA repair (HRR). Materials and Methods: We investigated Rad51 and HRR regulation in mouse embryo fibroblasts (MEFs) from PURA^{-/-} knockout mice that lack Pura. Results: Rad51 was induced in PURA^{-/-} MEFs but was repressed when Pura was ectopically expressed in these cells. Similarly Rad51 inversely correlated with the level of Pura in normal postnatal mouse brain. HIV-1 Tat stimulated HRR DNA repair of I-SceI induced DNA DSBs and the nuclear appearance of Rad51 foci. In contrast, Pura suppressed HRR DNA repair, Rad51 expression, and Rad51 foci formation. Conclusion: Tat stimulates the Rad51 promoter involving both Pura-dependent and Pura-independent mechanisms. Interaction between Pura and Tat may have opposing effects on Rad51 expression. The effects on HRR may contribute to HIV-1 associated pathogenesis.*

Pur α is a protein that was originally purified from mouse brain based on its ability to bind to a DNA sequence derived from the promoter of the mouse myelin basic protein gene (1, 2). Human Pur α was characterized by its ability to bind to a DNA sequence present upstream of the human c-Myc gene and was cloned from HeLa cells (3, 4). The sequence of mouse Pur α (5) is almost identical to human Pur α (4) with only two amino acid residues differing. The DNA-binding

domain of Pur α is strongly conserved throughout evolution. Pur α is a member of the Pur family of proteins along with Pur β (4) and Pur γ which has two isoforms that arise from the usage of alternative polyadenylation sites (6), and it is expressed in virtually every metazoan tissue (7). Pur α is a multifunctional protein that can bind to both DNA and RNA and functions in the initiation of DNA replication, control of transcription and mRNA translation (7, 8). A novel role has recently been described for Pur α in the transport and targeting of mRNAs in neurons (9, 10).

It is also becoming clear that Pur α is a major player in the regulation of the cell cycle and oncogenic transformation. Pur α binds to several cellular regulatory proteins including the retinoblastoma protein (11), E2F-1 (12, 13, 14), Sp1 (15) and YB-1 (16). Levels of Pur α fluctuate during the cell cycle, declining at the onset of S-phase and peaking during mitosis (17). Microinjection of Pur α into NIH-3T3 cells causes cell cycle arrest at either the G1/S or G2/M checkpoints (18). Expression of Pur α in Ras-transformed NIH-3T3 cells inhibits their ability to grow in soft agar (19). Ectopic overexpression of Pur α suppresses the growth of several transformed and tumor cells including glioblastomas (20). Gene expression profiling in chronic myeloid leukemia patients revealed a down-regulation of Pur α expression (21) and deletions of Pur α have been reported in myelodysplastic syndrome, a condition that can progress to acute myelogenous leukemia consistent with a role for Pur α as a tumor suppressor (22).

The generation and analysis of knockout mice with targeted inactivation of the PURA gene that encodes Pur α revealed that Pur α has an essential role in postnatal brain development (23). Mice with targeted disruption of the PURA gene in both alleles (PURA^{-/-}) appear normal at birth, but at two weeks of age, develop neurological abnormalities culminating in death by four weeks. There are much fewer cells in the brain cortex, hippocampus and cerebellum of

Correspondence to: Kamel Khalili, 1900 North 12th Street, 015-96, Room 203, Philadelphia, PA 19122, U.S.A. Tel: +215 204 0678, Fax: +215 204 0679, e-mail: kamel.khalili@temple.edu

Key Words: Pur-alpha, HIV-1 Tat, Rad51, DNA repair.

PURA^{-/-} mice than in PURA^{+/+} controls due to a lack of proliferation of precursor cells in this region. This implicates Purα in the regulation of developmentally timed DNA replication in specific cell types in the brain. These knockout mice also provide a source of mouse embryo fibroblast primary cultures (MEFs) with PURA deletion that can be used in comparison with MEF cells as an experimental system to examine the cellular functions of Purα.

It has been found that Purα has a role in the life cycle of certain viruses including JC virus (JCV) and human immunodeficiency virus type-1 (HIV-1). In this regard, some viral regulatory proteins target Purα and these include the Tat transactivator protein of the HIV-1 (24) and the large T-antigen of the human neurotropic polyomavirus JC (25). HIV-1 Tat protein transactivates the JCV late promoter (26) and this transactivation is mediated by the interaction of Purα with Tat which stimulates JCV late promoter transcription (27) and JCV DNA replication (28). Purα-mediated activation of JCV by Tat may be important in the pathogenesis of progressive multifocal leukoencephalopathy, which is caused by JCV and occurs predominantly in individuals with HIV-1/AIDS (29). In the life cycle of HIV-1 itself, Purα has recently been found to have a role in the Rev-mediated nuclear export of certain HIV-1 mRNAs (30).

Interestingly, we have found that both Purα and HIV-1 Tat affect cellular DNA repair, specifically homologous recombination-directed DNA repair (HRR) of double-strand breaks (DSBs). Purα negatively affects HRR and represses the expression of the HRR protein Rad51 (31) whereas Tat increases HRR DSB repair activity and induces the expression of Rad51 (32). In light of these findings and our earlier observations that Tat and Purα physically and functionally interact, we examined the effect of Tat and Purα on Rad51 and HRR utilizing MEFs derived from the PURA^{-/-} mice and PURA^{+/+} controls.

Materials and Methods

Cell lines and cell cultures. Primary human fetal astrocytes were cultured from human fetal tissues as described previously (33, 34). Primary mouse embryo fibroblasts (MEFs) were derived from homozygous PURA^{-/-} mouse embryos, heterozygous PURA^{+/-} mouse embryos and their partner wild-type (WT) PURA^{-/-} mouse embryos. PURA^{-/+}(+Purα) MEFs were established by transfecting PURA^{-/-} MEFs with pcDNA3-Purα containing full length Purα cDNA and PURA^{-/-} MEFs/GFP-Purα with pTRE-GFP-Purα containing tagged GFP Purα. MEFs were cultured in DMEM medium supplemented with 10% FBS and antibiotics. Cells were maintained in a humid incubator at 37°C with 5% CO₂.

Immunofluorescence. Cells were plated for immunofluorescence microscopy and grown on poly-D-lysine coated glass chamber slides. Spontaneous Rad51 foci formation was determined in exponentially growing cells. DNA damage-induced Rad51 foci were determined in cells treated with 2 mM HU for 24 hours.

Cells were grown for 24 h in chamber slides and then treated with 2 mM hydroxyurea (HU) for another 24 hours. Untreated cells and treated cells were fixed for 15 minutes in 3% paraformaldehyde at room temperature. After a wash in phosphate buffered saline (PBS), cells were permeabilized for 15 min at room temperature in PBS containing 0.3% Triton X-100. Slides were washed three times in PBS for 5 min and subsequently blocked with PBS containing 10% normal goat serum for 30 min. Subsequently, anti-Rad51 polyclonal antibody was applied to the slides for 60 min at room temperature followed by a goat anti-rabbit secondary antibody conjugated with Rhodamine (1: 400 dilution, Molecular Probes, Eugene USA). After extensive washing, slides were mounted with coverslips using antifading solution containing DAPI. For detection of RAD51 foci, fluorescence microscopy was used. For each data point, at least 100 nuclei were analyzed. Nuclei containing more than 10 strongly fluorescing foci were counted as positive. Images of Rad51 foci were recorded using an inverted fluorescence microscopy with a digital deconvolution system with a 100x oil immersion objective. Thirty-two optical sections per nucleus were generated at 0.4 μm distance. The 3D staining cells were viewed and each fluorescence was counted using Slidebook software. The volume of Rad51 foci and GFP-Purα staining were quantified in the same total volume of nucleus. The colocalization volume is counted by merging the red for Rad51 and green labeling for Purα.

Rad51 promoter assay by transient transfection. Cells were transfected with pRad51-LUC, which contains the human Rad51 promoter (-722 to +37) driving expression of the luciferase gene (35). After 48 hours, the cells were lysed and luciferase activity was measured using a luciferase assay kit according to the manufacturer's instructions (Promega, Madison, WI, USA).

Homologous recombination assay. pDR-GFP (obtained from Dr. Jasin's laboratory, Sloan-Kettering Institute, New York, NY, USA) containing a mutated GFP gene with an 18 bp *I-SceI* site was transfected into PURA^{-/-} MEFs cells using Lipofectamine 2000 (Invitrogen Corporation, Carlsbad, CA). The stable transfected cell lines were selected by growing in medium containing 5 μg/ml of puromycin. Puromycin-resistant colonies were screened by Southern blot analyses for an intact DR-GFP reporter. A 714-bp GFP coding fragment obtained by PCR system 9700 (Perkin Elmer Inc., Wellesley, MA) with primers forward 5'-ATGGTGAGC AAGGGCGAGGAGCT-3' and reverse 5'-CTTGTACAGCTCG TCATGCCGA-3' from template pDR-GFP labeled with ³²P as the probe. Ten micrograms of genomic DNA from puromycin-resistant colonies were digested by *SalI* and *HindIII* and separated on a 0.7% agarose gel and transferred to a nylon membrane. Hybridization was carried out under standard conditions using the ³²P-labeled 714-bp probe to test whether these puromycin-resistant colonies have integrated an intact DR-GFP fragment. A radiosensitive screen exposed with the hybridized membrane was analyzed using a Phosphorimager (Storm 840; Amersham) with the ImageQuant analysis software (Amersham). To evaluate HRR of DNA DSBs, MEF-DRGFP cells were prepared for transfection by pCMV3xnlS-I-SceI DNA using LipofectaminePlus as described above. At 48 h post transfection, cells were collected by trypsinization and GFP expression assayed by flow cytometry (Beckman Coulter, Epics Elite ESP) using an argon ion laser emitting at 488 nm. The frequency of recombination events was calculated from the

frequency of GFP-positive cells. All experiments were repeated at least three times and the results are shown as average.

Cell cycle analysis. To analyze cell cycle profiles, harvested cells were labeled for flow cytometer with propidium iodide solution. Cell cycle was measured by a FACScan flow cytometry and analyzed by using CELLQUEST software.

Results

Pura regulates Rad51 expression. Rad51 plays a critical role in repairing DNA double-strand breaks by promoting homologous recombination-directed pairing and strand exchange between damaged DNA and homologous DNA duplexes. It has been shown that Rad51 is down regulated in primary cells and overexpressed in proliferating cells and tumor tissues. In contrast, our previous studies have shown that Pura expression is increased in brain tissue during development, reaching a peak stage at postnatal day 15 (23). In this study, we found that Rad51 and PCNA levels were reduced at day 15, the peak of Pura gene expression (Figure 1A). Rad51 expression was increased in *PURA*^{+/-} and *PURA*^{-/-} mouse embryo fibroblasts (MEFs) (Figure 1B) relative to wild-type mice. Introduction of ectopically expressed Pura reduced the level of Rad51 in *PURA*^{-/-} MEFs. These results suggest that Pura downregulates Rad51 gene expression or promotes Rad51 protein degradation.

It is well established that locations where DNA damage has been induced and stalled DNA replication forks recruit Rad51 at sites of DSB repair, leading to the formation of foci visible by immunofluorescence (IF) using anti-Rad51 antibodies (Figure 2A). Rad51 foci seen in untreated cells are thought to reflect spontaneous DSB or DSB generated during DNA replication from other forms of damage. To investigate whether increased Rad51 expression in *PURA*^{-/-} cells reflects increased DNA damage or repair activity, we examined Rad51 foci formation. The results in Figure 2B indicate that there are more Rad51 foci positive cells in untreated *PURA*^{-/-} cells (*Pura*⁻) cells than wild-type cells (statistical significant in t-test $p < 0.05$; Figure 2B left) and that ectopic expression of Pura in *PURA*^{-/-} cells (+*Pura*) reduces the number of Rad51 foci positive cells to the levels seen in wild-type cells. Long-term exposure to the DNA replication inhibitor, hydroxyurea (HU), causes stalled replication forks to collapse into double-strand breaks. We found that the percent of cells with Rad51 foci increases in wild-type, *PURA*^{-/-} and *PURA*^{-/-} (+*Pura*) following treatment with HU for 24 hours (Figure 2B, right). A slightly higher percentage of cells containing Rad51 foci were found in *PURA*^{-/-} cells than in wild-type and *PURA*^{-/-} (+*Pura*) cells. Furthermore, we found that Pura is translocated into the nucleus in some of HU-treated *PURA*^{-/-} (+*Pura*) cells (Figure 2C). There are significantly less Rad51 foci in Pura positive nuclei compared to Pura negative nuclei (Figure




Figure 1. *Pura* suppresses Rad51 expression. A. The expression of Rad51 and Pura in the brain was measured during mouse development. Whole tissue protein extracts from mouse brains were prepared at postnatal days 5, 10 & 15 and were immunoblotted with antibodies against Pura, PCNA and Rad51. Grb-2 protein was used as a loading control. B. Expression of Rad51 was measured in MEFs isolated from *PURA*^{+/+} (wild-type), *PUR*^{+/-} and *PUR*^{-/-} mice. The lane labeled +*Pura* represents *PUR*^{-/-} MEFs where Pura was ectopically expressed as described in Materials and Methods.

2D). These results demonstrate that Pura significantly reduced Rad51 foci and Rad51 expression and that these events are related to both spontaneous DNA damage (-HU) and replication stress (+HU).

Pura and HIV-1 Tat regulate Rad51 transcription. Promoter analysis showed that there are several Pura binding sites within the Rad51 promoter region. To test whether Pura regulates Rad51 promoter activity, we used a reporter plasmid with the Rad51 promoter driving luciferase expression in wild-type, *PURA*^{-/-} and *PURA*^{-/-} (+*Pura*-) MEFs. As shown in Figure 3A, we found that HIV-1 Tat significantly stimulated Rad51 promoter activity.

Pura and HIV-1 Tat interfere with homologous recombination-directed DNA Repair (HRR). To analyze the effect of Pura on HRR, *PURA*^{-/-} and *PURA*^{-/-} (+*Pura*) cells were transfected with a recombination substrate DR-GFP that carries a tandem repeat of the GFP gene in which one copy is inactivated by the I-SceI sequence (scGFP) and the other by truncation of the C- and N-termini (iGFP). HRR of




Figure 2. *Purα* suppresses *Rad51* foci formation. *Rad51* foci formation was visualized in MEFs isolated from *PURA*^{+/+}, *PUR*^{-/-} and *PUR*^{-/-} MEFs expressing *Purα*. Cells were treated with and without 2 mM hydroxyurea for 24 hours, labeled with anti-*Rad51* antibody and visualized under fluorescence microscopy. Red represents rhodamine-labeling of *Rad51* foci and blue is DAPI labeling of nuclear DNA. Cells containing more than 10 *Rad51* foci were counted as positive. **A.** A typical *Rad51* foci labeling from untreated *PUR*^{-/-} MEFs. **B.** The plot of percentage of *Rad51* foci positive cells for the three types of cells either untreated (left histogram) or treated with 2 mM HU for 16 hours. **C.** *PURA*^{-/-} MEFs ectopically expressing *Purα* and treated with 2 mM HU for 16 hours. **D.** The plot of percentage of *Rad51* foci in HU-treated *PURA*^{-/-} (+*Purα*) cell nuclei that are either immunopositive for *Purα* (left) or immunopositive (right). The means (symbols) and standard errors (error bars) were derived from three independent experiments.

the I-SceI induced DSB restores the GFP gene. Cell lines that have stably integrated the construct were obtained by puromycin selection. To measure the efficiency of HRR, the pCMV3nls-I-SceI plasmid was transfected into these cells and the fraction of GFP expressing cells was measured by




Figure 3. *HIV-1 Tat* stimulates *Rad51* promoter activity and homologous recombination. **A.** MEFs were transfected with pGL3*Rad51*-luciferase plasmid along with *pcDNA3*-vector, *pcDNA3*-*Purα* or *pcDNA*-*Tat* expression plasmid using Lipofectamine (Invitrogen). Luciferase activity was determined after 36 h. The data represent results from three independent experiments. **B.** The effects of *HIV-1 Tat* and *Purα* on homologous recombination-directed DNA repair were examined. *PURA*^{-/-} and *PURA*^{-/-} (+*Purα*) MEFs were transfected with *pcDNA3* or *pcDNA3*-*TAT* along with pCMV-3nls-I-SceI for 48 hours. The effect of *HIV-1 TAT* in *PURA*^{-/-} (+*Purα*) cells was significantly different from control, **p*<0.01, but that in *PURA*^{-/-} cells was not, ***p*>0.05.

flow cytometry 48h later. *Purα* expression reduces by half the number of GFP positive cells observed after I-SceI transfection. These results suggest that *Purα* inhibits HRR.

Previous studies have demonstrated that *HIV-1 Tat* increases *Rad51* expression and enhances homologous recombination (32). Further, *HIV-1 Tat* has been shown to interact with *Purα* and this interaction is involved in the stimulation of transcription of the JC virus promoter by *Tat* (27). To further investigate the effect of *HIV-1 Tat* on HRR and the role of *Purα*, we expressed *HIV-1 Tat* in *PURA*^{-/-}




Figure 4. HIV-1 Tat activates primary human fetal astrocytes. Cells were grown for one week before transfected with pcDNA3 HIV-1 Tat plasmid DNA. 48 hours after transfection, cells were labeled with rabbit anti-HIV-1 Tat and with mouse anti-BrdU, anti-PCNA or anti-Rad51 antibodies. Labeled cells were analyzed by immunofluorescence microscopy. Top row, astrocytes expressing Tat are actively replicating DNA (BrdU-positive, top panels). In Tat-positive mitotic cells, PCNA (middle panels) and Rad51 (lower panels) are expressed between the chromosomes.

cells in the presence or absence of co-expressed Pura. We found that HIV-1 Tat expression significantly increased HRR in *PURA*^{-/-} cells and that in the presence of co-expressed Pura, Tat enhanced HRR significantly over cells expressing Pura alone (Figure 3B). These results suggest that Pura and HIV-1 Tat have opposite effects on HRR.

HIV-1 Tat induces DNA replication and Rad51 expression in primary human fetal astrocytes. Our previous studies have shown that Rad51 expression is activated in PML tissue from patients infected with HIV-1 and JCV (36). To determine whether HIV-1 Tat affects Rad51 expression in primary cells, we transfected primary human fetal astrocytes with pCMV-

Tat construct. Figure 4 shows that DNA replication and Rad51 expression are activated in HIV-1 Tat expressing cells. Furthermore, HIV-1 Tat positive cells entered mitosis and are particularly arrested at anaphase. These findings indicate that HIV-1 Tat may induce unexpected DNA replication and mitosis, possibly leading to chromosomal instability.

Discussion

Our results indicate that Pura inhibits the level of expression of Rad51 protein and the formation of Rad51 foci. Thus, the Pura protein levels in the developing mouse brain and in cultured MEFs are inversely correlated with the level of

Rad51. The decreased levels of Rad51 may be caused by either the inhibition of Rad51 transcription or an increase in the rate of Rad51 protein degradation by Pura. There are a number of possible reasons for the increased incidence of Rad51 foci formation in the absence of Pura. During DNA replication, DNA replication arrest can induce DNA damage and unrepaired single-strand breaks can transit into double-strand breaks (DSBs). It is possible that Pura acts as a caretaker protein that is involved in the stabilization and repair of stalled replication forks and/or DSBs. In the absence of Pura, this would lead to an increase in DSBs and enhanced Rad51 foci formation. Our earlier studies on the involvement of Pura in the cellular response to DNA replication stress support this hypothesis (31).

Rad51 becomes recruited to sites of DNA damage and binds to single strand DNA to form bright foci. Rad51 is involved in homologous recombination-directed DNA repair (HRR), which provides an efficient and faithful pathway of repairing DNA damage, especially during DNA replication. However, as discussed in the introduction, Pura has the properties of a tumor suppressor protein and ablation of its expression in certain cancers is associated with genetic instability. Thus it could be possible that loss of the DSB caretaker function of Pura is responsible for this observation. Alternatively, Pura is involved in cell cycle arrest (18) and in the absence of Pura, it is possible that cell cycle progression may occur inappropriately before DNA repair has occurred. A third possibility is that the induction of Rad51 in the absence of Pura is responsible. Paradoxically, when Rad51 is overexpressed, it reduces DSB-induced HRR DNA repair perhaps because excess Rad51 blocks DNA strand exchange (37). Indeed, the aberrant increase in Rad51 expression that is found in tumor cells may contribute to genomic instability by promoting abnormal recombination (38, 39, 40). In another recent study, we reported that JC virus infection promotes genomic instability that is associated with a large induction of Rad51 protein level (36).

Another protein that can modulate the expression of Rad51 and HRR is HIV-1 Tat as we have recently reported (32). Unlike Pura, Tat has a positive effect, *i.e.*, it increases the level of Rad51 and the rate of HRR. Pura and Tat are able to physically and functionally interact and this is involved in regulating the transcription of certain genes, *e.g.*, the viral late promoter of JCV (27) and the cellular TGF β -1 promoter (41). In the light of these findings, we have now examined the effect of Pura and Tat interaction on Rad51 and HRR. Interestingly, Tat stimulated the Rad51 promoter in the absence of Pura (Figure 3A, column 8) but this induction was significantly enhanced in the presence of Pura (Figure 3A, column 9). Similarly, HRR was stimulated by Tat both in the presence and in the absence of Pura (Figure 3B). This suggests that the regulation of Rad51 and HRR by Tat involves both Pura-dependent and Pura-independent mechanisms. With regard to the Pura-independent mechanism of regulation, we have

recently found that wild-type p53 negatively regulates the Rad51 promoter but the T-antigen of JCV, which antagonizes p53 function, reverses this inhibition (35). Since reciprocal modulations also occur between p53 and Tat (42), it is possible that the Pura-independent mechanism of Rad51 promoter regulation involves p53.

In conclusion, events involving the effects of Pura and HIV-1 Tat on Rad51 expression and HRR may be important in understanding HIV-1 associated pathogenesis.

Acknowledgements

We thank past and present members of the Center for Neurovirology for their insightful discussion and sharing of ideas and reagents. We also wish to thank C. Schriver for editorial assistance. This work was supported by grants awarded by the NIH to KK.

References

- Haas S, Gordon J and Khalili K: A developmentally regulated DNA-binding protein from mouse brain stimulates myelin basic protein gene expression. *Mol Cell Biol* 13: 3103-3112, 1993.
- Haas S, Thatikunta P, Steplewski A, Johnson EM, Khalili K and Amini S: A 39-kD DNA-binding protein from mouse brain stimulates transcription of myelin basic protein gene in oligodendrocytic cells. *J Cell Biol* 130: 1171-1179, 1995.
- Bergemann AD and Johnson EM: The HeLa Pur factor binds single-stranded DNA at a specific element conserved in gene flanking regions and origins of DNA replication. *Mol Cell Biol* 12: 1257-1265, 1992.
- Bergemann AD, Ma ZW and Johnson EM: Sequence of cDNA comprising the human pur gene and sequence-specific single-stranded DNA-binding properties of the encoded protein. *Mol Cell Biol* 12: 5673-5682, 1992.
- Ma ZW, Bergemann AD and Johnson EM: Conservation in human and mouse Pur alpha of a motif common to several proteins involved in initiation of DNA replication. *Gene* 149: 311-314, 1994.
- Liu H and Johnson EM: Distinct proteins encoded by alternative transcripts of the PURG gene, located contrapodal to WRN on chromosome 8, determined by differential termination/polyadenylation. *Nucleic Acids Res* 30: 2417-2426, 2002.
- Johnson EM: The Pur protein family: clues to function from recent studies on cancer and AIDS. *Anticancer Res* 23: 2093-2100, 2003.
- Gallia GL, Johnson EM and Khalili K: Puralpha: a multifunctional single-stranded DNA- and RNA-binding protein. *Nucleic Acids Res* 28: 3197-3205, 2000.
- Johnson EM, Kinoshita Y, Weinreb DB, Wortman MJ, Simon R, Khalili K, Winckler B and Gordon J: Role of Pur alpha in targeting mRNA to sites of translation in hippocampal neuronal dendrites. *J Neurosci Res* 83: 929-943, 2006.
- Kanai Y, Dohmae N and Hirokawa N: Kinesin transports RNA: isolation and characterization of an RNA-transporting granule. *Neuron* 43: 513-525, 2004.
- Johnson EM, Chen PL, Krachmarov CP, Barr SM, Kanovsky M, Ma ZW and Lee WH: Association of human Pur alpha with the retinoblastoma protein, Rb, regulates binding to the single-stranded DNA Pur alpha recognition element. *J Biol Chem* 270: 24352-24360, 1995.

- 12 Darbinian N, Gallia GL, Kundu M, Shcherbik N, Tretiakova A, Giordano A and Khalili K: Association of Pur alpha and E2F-1 suppresses transcriptional activity of E2F-1. *Oncogene* 18: 6398-6402, 1999
- 13 Darbinian N, White MK, Gallia GL, Amini S, Rappaport J and Khalili K: Interaction between the Pura and E2F-1 transcription factors. *Anticancer Res* 24: 2585-2594, 2004.
- 14 Darbinian N, White MK and Khalili K: Regulation of the Pur-alpha promoter by E2F-1. *J Cell Biochem* 99: 1052-1063, 2006.
- 15 Tretiakova A, Steplewski A, Johnson EM, Khalili K and Amini S: Regulation of myelin basic protein gene transcription by Sp1 and Puralpha: evidence for association of Sp1 and Puralpha in brain. *J Cell Physiol* 181: 160-168, 1999.
- 16 Safak M, Gallia GL and Khalili K: Reciprocal interaction between two cellular proteins, Puralpha and YB-1, modulates transcriptional activity of JCVCY in glial cells. *Mol Cell Biol* 19: 2712-2723, 1999.
- 17 Itoh H, Wortman MJ, Kanovsky M, Uson RR, Gordon RE, Alfano N and Johnson EM: Alterations in Pur(alpha) levels and intracellular localization in the CV-1 cell cycle. *Cell Growth Differ* 9: 651-665, 1998.
- 18 Stacey DW, Hitomi M, Kanovsky M, Gan L and Johnson EM: Cell cycle arrest and morphological alterations following microinjection of NIH3T3 cells with Pur alpha. *Oncogene* 18: 4254-4261, 1999.
- 19 Barr SM and Johnson EM: Ras-induced colony formation and anchorage-independent growth inhibited by elevated expression of Puralpha in NIH3T3 cells. *J Cell Biochem* 81: 621-638, 2001.
- 20 Darbinian N, Gallia GL, King J, Del Valle L, Johnson EM and Khalili K: Growth inhibition of glioblastoma cells by human Pur(alpha). *J Cell Physiol* 189: 334-340, 2001.
- 21 Bruchova H, Borovanova T, Klamova H and Brdicka R: Gene expression profiling in chronic myeloid leukemia patients treated with hydroxyurea. *Leuk Lymphoma* 43: 1289-1295, 2002.
- 22 Lezon-Geyda K, Najfeld V and Johnson EM: Deletions of PURA, at 5q31, and PURB, at 7p13, in myelodysplastic syndrome and progression to acute myelogenous leukemia. *Leukemia* 15: 954-962, 2001.
- 23 Khalili K, Del Valle L, Muralidharan V, Gault WJ, Darbinian N, Otte J, Meier E, Johnson EM, Daniel DC, Kinoshita Y, Amini S and Gordon J: Puralpha is essential for postnatal brain development and developmentally coupled cellular proliferation as revealed by genetic inactivation in the mouse. *Mol Cell Biol* 23: 6857-6875, 2003.
- 24 Gallia GL, Darbinian N, Tretiakova A, Ansari SA, Rappaport J, Brady J, Wortman MJ, Johnson EM and Khalili K: Association of HIV-1 Tat with the cellular protein, Puralpha, is mediated by RNA. *Proc Natl Acad Sci USA* 96: 11572-11577, 1999.
- 25 Gallia GL, Safak M and Khalili K: Interaction of the single-stranded DNA-binding protein Puralpha with the human polyomavirus JC virus early protein T-antigen. *J Biol Chem* 273: 32662-32669, 1998.
- 26 Tada H, Rappaport J, Lashgari M, Amini S, Wong-Staal F and Khalili K: Trans-activation of the JC virus late promoter by the tat protein of type 1 human immunodeficiency virus in glial cells. *Proc Natl Acad Sci USA* 87: 3479-3483, 1990.
- 27 Krachmarov CP, Chepenik LG, Barr-Vagell S, Khalili K and Johnson EM: Activation of the JC virus Tat-responsive transcriptional control element by association of the Tat protein of human immunodeficiency virus 1 with cellular protein Pur alpha. *Proc Natl Acad Sci USA* 93: 14112-14117, 1996.
- 28 Daniel DC, Kinoshita Y, Khan MA, Valle LD, Khalili K, Rappaport J and Johnson EM: Internalization of exogenous human immunodeficiency virus-1 protein, Tat, by KG-1 oligodendrogloma cells followed by stimulation of DNA replication initiated at the JC virus origin. *DNA Cell Biol* 23: 858-867, 2004.
- 29 Khalili K, Gordon J and White MK: The polyomavirus, JCV, and its involvement in human disease. *Adv Exp Med Biol* 577: 274-287, 2006.
- 30 Kaminski R, Darbinian N, Sawaya BE, Slonina D, Amini S, Johnson EM, Rappaport J, Khalili K and Darbinian A: Puralpha as a cellular co-factor of Rev/RRE-mediated expression of HIV-1 intron-containing mRNA. *J Cell Biochem* 103: 1231-1245, 2008.
- 31 Wang H, Wang M, Reiss K, Darbinian-Sarkissian N, Johnson EM, Iliakis G, Amini S, Khalili K and Rappaport J: Evidence for the involvement of Puralpha in response to DNA replication stress. *Cancer Biol Ther* 6: 596-602, 2007.
- 32 Chipitsyna G, Slonina D, Siddiqui K, Peruzzi F, Skorski T, Reiss K, Sawaya BE, Khalili K and Amini S: HIV-1 Tat increases cell survival in response to cisplatin by stimulating Rad51 gene expression. *Oncogene* 23: 2664-2671, 2004.
- 33 Radhakrishnan S, Otte J, Enam S, Del Valle L, Khalili K and Gordon J: JC virus-induced changes in cellular gene expression in primary human astrocytes. *J Virol* 77: 10638-10644, 2003.
- 34 Radhakrishnan S, Gordon J, Del Valle L, Cui J and Khalili K: Intracellular approach for blocking JC virus gene expression by using RNA interference during viral infection. *J Virol* 78: 7264-7269, 2004.
- 35 White MK, Skowronska A, Gordon J, Del Valle L, Deshmane SL, Giordano A, Croul S and Khalili K: Analysis of a mutant p53 protein arising in a medulloblastoma from a mouse transgenic for the JC virus early region. *Anticancer Res* 26: 4079-4092, 2006.
- 36 Darbinian A, White MK, Akan S, Radhakrishnan S, Del Valle L, Amini S and Khalili K: Alterations of DNA damage repair pathways resulting from JCV infection. *Virology* 364: 73-86, 2007.
- 37 Kim PM, Allen C, Wagener BM, Shen Z and Nickoloff JA: Overexpression of human RAD51 and RAD52 reduces double-strand break-induced homologous recombination in mammalian cells. *Nucl Acids Res* 29: 4352-4360, 2001.
- 38 Raderschall E, Stout K, Freier S, Suckow V, Schweiger S and Haaf T: Elevated levels of Rad51 recombination protein in tumor cells. *Cancer Res* 62: 219-225, 2002.
- 39 Richardson C, Stark JM, Ommundsen M and Jasin M: Rad51 overexpression promotes alternative double-strand break repair pathways and genome instability. *Oncogene* 23: 546-553, 2004.
- 40 Xia SJ, Shammass MA and Shmookler Reis RJ: Elevated recombination in immortal human cells is mediated by HsRAD51 recombinase. *Mol Cell Biol* 17: 7151-7158, 1997.
- 41 Thatikunta P, Sawaya BE, Denisova L, Cole C, Yusibova G, Johnson EM, Khalili K and Amini S: Identification of a cellular protein that binds to Tat-responsive element of TGF beta-1 promoter in glial cells. *J Cell Biochem* 67: 466-477, 1997.
- 42 Li CJ, Wang C, Friedman DJ and Pardee AB: Reciprocal modulations between p53 and Tat of human immunodeficiency virus type 1. *Proc Natl Acad Sci USA* 92: 5461-5464, 1995.

Received February 12, 2008
Accepted February 28, 2008