

Review

## Molecular Biomarkers and Classification Models in the Evaluation of the Prognosis of Colorectal Cancer

MICHAIL SIDERIS and SAVVAS PAPAGRIGORIADIS

*Department of Colorectal Surgery King's College Hospital NHS Foundation Trust, Denmark Hill, London, U.K.*

**Abstract.** *Colorectal cancer (CRC) is the second leading cause of cancer-related death. Despite the progress that has been made towards the identification of the molecular*

*mechanisms involved in CRC, currently there are many unclear points. The current opinion is that microsatellite instability (MSI), CpG island methylator phenotype (CIMP) and chromosomal instability (CIN) seem to play a significant role. MSI is related to point mutations in defect mismatch repair system of DNA. There are two well-established MSI phenotypes: MSI-high (MSI-H) and MSI-low (MSI-L or MSS). CIN refers to a different cellular event which originates from the presence of an abnormal chromosome complement or number. CIMP is the third most commonly involved event, and is defined by widespread methylation of CpG islands of suppressor promoters, with two phenotypes: CIMP-high and CIMP-low which interact with MSI or CIN status V-raf murine sarcoma viral oncogene homolog B (BRAF) is a serine-threonine protein kinase that acts as a downstream effector of the Kirsten rat sarcoma viral oncogene homolog (KRAS) pathway. Various studies have revealed that BRAF V600E mutations appear to be a valid indicator of poor prognosis. KRAS is a proto-oncogene which encodes a GTP-ase involved in cellular response to extracellular stimuli. Its prognostic value is still controversial. However, wild-type KRAS is associated with better response to Endothelial Growth Factor Receptor inhibitors combined with standard chemotherapy. Loss of Heterozygosity, especially involving 18q, is a well-known potential mechanism for tumorigenesis that has been studied in CRC. Vascular endothelial growth factor is a pro-angiogenic factor linked with the aggressiveness of CRC. Emerging data show that cyclooxygenase 2 overexpression is significantly associated with worse outcomes in CRC. Recent studies highlight mi-croRNAs as promising prognostic biomarkers. More specifically, the down-regulation of miR-451, miR-625, miR-29c, miR-126, miR-129 and miR133 is purported to be a poor prognostic factor, while miR-224 was overexpressed in CRC.*

This article is freely accessible online.

**Abbreviations:** 5FU: 5 Fluorouracil; APC: adenomatous polyposis coli; BMI: body mass index; BRAF: v-raf murine sarcoma viral oncogene homolog B; BRAF V600E: v-raf murine sarcoma viral oncogene homolog B V600E mutation; CABLE1: CDK5 And ABL1 enzyme substrate; CIMP: CpG island Methylator Phenotype; CIN: chromosomal instability; COX-2: cyclooxygenase-2; CRC: colorectal cancer; DCC: deleted in colorectal carcinoma; DFS: disease-free survival; EGFR: epidermal growth factor; JCVT: John Cunningham (JC) virus T antigen; KRAS: kirsten rat sarcoma viral oncogene homolog; LINE-1 methylation: long interspersed nucleotide element-1 methylation; LOH: loss of heterozygosity; MAPK: mitogen-activated protein kinase; MLH-1: mutL homolog 1; MMR: mismatch repair (system); MSH2: mutS homolog 2; MSH6: mutS homolog 2; MSI: microsatellite instability; MSI-H: microsatellite instability – high; MSI-L: microsatellite instability – low; MSS: microsatellite stable; NRAS: neuroblastoma RAS viral (v-ras) oncogene homolog; PETACC: Pan-European Trials in Adjuvant Colon Cancer; PHLPP1: PH domain and leucine rich repeat protein phosphatase 1; PHLPP2: PH domain and leucine rich repeat protein phosphatase 2; PI3K/AKT: phosphoinositide-3-kinase / v-akt murine thy-moma viral oncogene; PI3KCA: phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha; PMS2: post-meiotic segregation increased 2 (S. cerevisiae); RFFL: ring finger and FYVE-like domain containing E3 ubiquitin protein ligase; SMAD2: mothers against decapentaplegic homolog 2; SMAD4: mothers against decapentaplegic homolog 4 (Drosophila); UFT: tegafur-uracil oral chemotherapy; VEGF: vascular endothelial growth factor.

**Correspondence to:** Dr. Michail Sideris, Department of Colorectal Surgery, Hambleton Wing, 3rd Floor, King's College Hospital NHS Foundation Trust, Denmark Hill, London, SE5 9RS, U.K. Tel: +44 20 3299 4667, e-mail: msideris@nhs.net

**Key Words:** CRC biomarkers, CIMP, CIN, MSI, KRAS, BRAF, miRNA, review.

Currently, treatment strategies and clinical outcomes in colorectal cancer (CRC) are determined by cancer stage as defined by TNM or relevant staging protocols (1), based on local tumor penetration, and spread to lymph nodes or other organs. The cornerstone of non-metastatic CRC treatment remains surgical resection. In patients with higher stage disease (III or high risk II) however, there is a significant risk of recurrence. Although certain histological factors such as tumor differentiation, grade, and lymphovascular invasion, have been identified as being to higher risk, there is still a lack of understanding of the molecular factors which may affect the risk of metastasis and recurrence.

In recent years, molecular biomarkers have generated interest as prognostic markers, or as in the case of Kirsten rat sarcoma viral oncogene homolog (*KRAS*), as factors involved in treatment choice. Generally, a biomarker is defined as a characteristic that can be objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes, or pharmacological response to a specified therapeutic intervention. Biomarkers can be either prognostic or predictive (2, 3). Prognostic values are defined by the estimated life expectancy post-diagnosis and treatment, whereas predictive biomarkers are related to the response of a patient to a relevant treatment strategy.

The purpose of a molecular classification is to identify similar characteristics among individual tumors, and then empirically predict the pathogenesis and biological behavior of a particular tumor (4, 5). The most accepted way of creating a classification model is to identify and correlate single cellular events that have been statistically proven to play a role in tumorigenesis.

The starting point of these efforts was the identification of significant stages in the progression from healthy epithelium to cancer (6, 7). Currently, some theories exist which explain the transition from adenoma, the first pre-cancerous type of lesion, to carcinoma and then invasion to other tissues. In this way, commonly involved events in CRC were identified and highlighted as the guiding points for a classification model. Subsequently, the first efforts to summarize and correlate these events were published, but none was actually adequately evidence-based to achieve translation to clinical practice.

Most of the current molecular classification models are based on Microsatellite Instability (MSI), Chromosomal Instability (CIN) and CpG island Methylator Phenotype (CIMP) and they correlate these event with other significant mutations *i.e.* *KRAS*, and *V-raf* murine sarcoma viral oncogene homolog B (*BRAF*) (7, 8).

In this review article, we summarize what is currently believed in terms of the popular potential biomarkers in CRC and set the parameters for further discussion and points of interest for further research.

## MSI

Microsatellite disorders are established as frequent events in CRC, occurring in around 22% of cases (9). There are many reports supporting the effect of MSI on CRC prognosis (3, 10-17).

Microsatellites are short repetitive DNA nucleotide sequences which are prone to frame-shift mutations and base-pair substitutions during DNA replication (3). They are involved in the DNA repair system and their mutations were initially associated with Lynch syndrome, which is an autosomal dominant disorder, highly associated with 3% of multiple CRC types. Mismatch Repair (*MMR*) genes involved in Lynch syndrome are mutS homolog 1 (*MLH1*), mutS homolog 2 (*MSH2*), mutS homolog 6 (*MSH6*) and post-meiotic segregation increased 2 (*PMS2*) (18). Loss of these genes results in defective MMR, which in turn results in MSI.

Sporadic CRC is more likely to be associated with *MLH1* mutation. In terms of MSI status classification, most studies divide it into three categories: MSI-high (MSI-H) at  $\geq 30\%$ , MSI-low (MSI-L) at 10-30% and microsatellite stable (MSS) (3, 18). Some other studies aim to simplify MSI status as being positive or negative (10, 19, 20). It has been questioned whether MSI-L exists or falls into the same category as MSS (21).

Most studies aiming to establish a classification model in CRC consider MSI status as one of the primary criteria for categorization (10, 20) (Table I). A recent multivariate analysis identified MSI alterations in 119/892 CRC samples and highlights MSI status as one of the major prognostic biomarkers (10). Another interesting study suggested a model which is mainly based on MSI status and its relationship with CpG island Methylator Phenotype (CIMP) and Chromosomal Instability (CIN). In the same study, it was noted that Adenomatous polyposis coli (*APC*) and *KRAS* mutations were less frequent in MSI tumors (13.3% and 10.9%, respectively) whereas *BRAF* V600E mutation is most often seen in MSI tumors (58.7%) (19).

There are many clinicopathological variables associated with MSI status. For instance, patients with MSI-H tumors in general have greater 5-year overall survival irrespective of stage compared to those with MSI-L and MSS tumors. The Pan-European Trials in Adjuvant Colon Cancer III (PETACC III) trial confirmed these retrospective findings and suggested that MSI-H is a strong prognostic factor for relapse-free and overall survival in patients with stage II and III CRC (3). Moreover, MSI-H tumors have been classified as proximally located, poorly-differentiated and with a higher incidence in female gender. They are also characterized by mucinous differentiation, increased age at onset, and round and vesicular nuclei with a prominent nucleolus (2, 10, 18-20).

However, contrary to the promising results in terms of the prognostic value of MSI status, no clear relationship between

Table I. *Baseline characteristics of MSI tumors. This table provides a summary of the baseline characteristics of MSI tumors and their correlation with KRAS and BRAF mutations.*

MSI	Summary of basic features
MSI-positive (high)	Present in around 22% of CRC cases (9, 10) Greater 5-year survival, proximal location, poor differentiation, female gender, mucinous differentiation (2, 3, 10, 18-20)
MSI in sporadic CRC	Association with <i>MLH1</i> (3,18)
MSI in LS	Association with <i>MLH1</i> , <i>MSH2</i> , <i>MSH6</i> , <i>PMS2</i> (18)
Correlation with <i>KRAS/BRAF</i> mutations	<i>BRAF</i> V600E is associated in 58.7% with MSI+ CRC (19) <i>KRAS</i> mutations present in 10.9% of MSI+ CRC (10)
Predictive value	Predictive value is still controversial (2, 20)

MLH-1: mutL homolog 1; MSH2: mutS homolog 2; MSH6: mutS homolog 2; PMS2: postmeiotic segregation increased 2.

MSI status and response to neoadjuvant chemotherapy has yet been proven. However, Ribic *et al.* concluded that MSI-H status can be associated with poor response to 5-Fluorouracil-based chemotherapy compared to MSI-L and MSS (2). Nevertheless, there are multiple studies with conflicting results in terms of the predictive value of MSI status and thus this is still under discussion (20).

### CIMP

Most studies develop the CRC classification model based on CIMP status and its correlation to CIN and MSI (4, 18-20, 22). CIMP is defined as hypermethylation of CpG island promoter. CIMP results in transcriptional silencing of specific tumor-suppressor and DNA repair genes, including *MLH1* (19). Some studies describe CIMP status as CIMP-high, CIMP-low and CIMP-negative (18). There is a discussion whether CIMP low and CIMP negative fall into the same category, resulting in two categories, CIMP-positive and CIMP-negative (22); we use this for simplicity here.

There have been some reasonable studies which associate MSI and CIMP status with CIN, resulting in a classification model which attempts to identify and link macroscopic variables to molecular features (19). In a recent study, Simons *et al.* suggest a classification based on MSI, CIMP and CIN. According to that, MSI and CIMP-positive tumors were more proximally located (85.7% and 51.9%, respectively), whereas CIMP-positive and CIN or CIN-only tumors were distally located (distal colon to rectum 52.3% and 82.1%, respectively). Triple-negative tumours were located in both proximal and distal colon and in 89% of them *BRAF V600E* mutation was identified. Regarding CIMP-only tumors, the average age at diagnosis was 67.6 years, 51.9% were located proximally, and the differentiation grade was II at the stage of data analysis. *BRAF V600E* mutation was present only in 18.5% and p53 overexpression was noted in 66.7% (19).

### CIN

Tumors with CIN have chromosomal gains and losses, with or without structural rearrangements, possibly reflecting an increased mutation rate; this is called aneuploidy and polyploidy, respectively (10, 23, 24), and occurs in around 60% of CRC cases (2). The mechanism underlying CIN is not yet understood. Most studies tend by definition to differentiate MSI-positive tumors from CIN, as CIN reflects a separate mechanism of carcinogenesis (19, 20). MSI-positive and CIN-positive tumors have been described but they tend to macroscopically appear more like MSI-positive (10). The overlap between CIN and MSI is not clear (19). CIN-positive tumors are generally associated with poor prognosis and tend to be well- or moderately differentiated (10). Simons *et al.* hold that CIN-only tumors are more frequent in men (54.7%), 76.5% at the stage of diagnosis are grade 2 and there is subsequent p53 overexpression (19). However, *BRAF V600E* mutation tends not to be involved in such a mechanism of tumorigenesis (10, 19), and some studies do link *KRAS* mutations with this molecular phenotype (19) (20). CIN tumors are located more often distally. CIMP- and CIN-positive tumors have been described (19).

Domingo *et al.* have described 3% of MSI-H and CIN-positive tumors tending to have characteristics closer to MSI-H CIN-negative tumors *i.e.* right-sided location and increased *BRAF* mutation, while p53 mutation and 17p LOH was not increased significantly, and thus there was no association (10).

### KRAS Mutation

*KRAS* is a proto-oncogene involved in cellular response to extracellular stimuli (3, 25). In cases of *KRAS* mutation, there is a structural activation of downstream signaling pathways *i.e.* mitogen-activated protein kinase (*MAPK*) and phosphoinositide-3-kinase/v-akt murine thymoma viral oncogen (*PI3K/AKT*) pathways (2), and through this mechanism, tumor cells are more

resistant to inhibition of surface receptors of tyrosine kinase such as epidermal growth factor receptor (EGFR). The MAPK pathway was recently found to be one of the fundamental mechanisms involved in tumorigenesis (9).

Point mutations in codons 12, 13 and 61 of *KRAS* gene are noted in about 30-40% of CRC (2) and in total, around 85 *KRAS* mutations have been identified (9). These different molecular phenotypes of *KRAS* may result in different pathways of carcinogenesis, which could alter the macroscopic phenotype.

However, it is still controversial whether *KRAS* can be considered as a valid biomarker or not, as recent studies suggest there are no clinicopathological feature to support this (3, 9). There are some controversial data emerging that link *KRAS* mutation with poor prognosis. Phipps *et al.* found *KRAS* mutation to be associated with poorer prognosis compared to wild-type *KRAS* (26). Most studies link *KRAS* mutation with certain molecular phenotypes *i.e.* CIN, whereas they are less likely to be noted in MSI-positive tumors (3, 10, 19, 20)

An interesting study in Moroccan patients with advanced CRC showed that 76.09% of patients had wild-type *KRAS* genotype, whereas 23.91% were *KRAS* mutants. The majority of *KRAS* mutations referred to an amino acid substitution of glycine by aspartic acid (68.2%) (27).

On the other hand, despite the debating prognostic value of *KRAS*, it has been established that wild-type *KRAS* is associated with better response to EGFR inhibitors in terms of adjuvant chemotherapy setting *i.e.* cetuximab (1, 3, 9, 28-31). An interesting study by Di Bartolomeo *et al.* in 2013 concluded that patients with *KRAS*/neuroblastoma RAS viral (v-ras) oncogene homolog, *BRAF* and *p53* wild-type tumors had the maximum benefit from treatment with cetuximab, oxaliplatin and tegafur-uracil oral chemotherapy (32) Nevertheless, Selcukbirik *et al.* found that *BRAF* and *KRAS* mutations do not seem to be potential biomarkers in the treatment of metastatic CRC with bevacizumab therapy (33). Bruera *et al.* also noted a worse prognosis in patients with metastatic CRC and *KRAS* (c35 G)/*BRAF* mutations (34), and a similar hypothesis is supported by Loupakis *et al.* where EGFR ligand was significantly modulated by cetuximab plus irinotecan therapy (35).

Tian *et al.* state that a combined signature of *KRAS/BRAF* and phosphatidyli-nositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha (*PI3KCA*) potentially offered an optimized source of information for cetuximab responsiveness of the tumor (36). Saridaki *et al.* concluded that *KRAS* and *BRAF* mutations, and EGFR expression can be used as biomarkers to further select patients undergoing anti-EGFR therapy (37).

## **BRAF Mutation**

*BRAF* gene encodes a serine-threonine kinase that acts as an inhibitor of the RAS/MAPK intracellular signaling pathway. Mutations of *BRAF* occur at an early stage of colorectal carcinogenesis (38).

In terms of correlations of *BRAF* with the main cellular events, most studies associate *BRAF* V600E activating mutation with sporadic MSI-H CRCs (2, 3, 9, 10, 19) and it is linked with poor prognosis (39) (Table II). What is more, no association between *BRAF* mutations and CIN has yet been noted (2, 10, 19). According to Doming *et al.*, *BRAF* alterations occur in 10% of CRC cases and there is a negative association with *KRAS* mutation (10), whereas there is primary positive association with MSI and CIMP-high (4, 31).

In a recent study by Lochhead *et al.*, *BRAF* mutation occurred in 10% to 20% of CRC and was associated with MSI-high through its relationship to CIMP-high (13). The same study concluded that *BRAF* mutations are associated with inferior prognosis. Eklof *et al.* analyzing the mutation status in *KRAS* and *BRAF*, and *PIK3CA* and *PTEN* expression in two separate CRC cohorts, state that *KRAS* and *BRAF* status are important in the establishment of the prognosis in CRC and should always be considered (40). Similarly, Yokota *et al.* conclude that mutated *BRAF* is one of the most powerful prognostic markers in CRC (41).

Nevertheless, *BRAF* as a predictive marker is still under discussion (3, 29). There are some studies which tend to use a signature of *BRAF/KRAS* as a predictive factor for the response to EGFR inhibitors (32-37). Another interesting meta-analysis (42), taking into consideration 21 trials including 5229 patients, concluded that *BRAF* mutation is a predictive biomarker for poor prognosis in patients with metastatic CRC undergoing therapy with anti-EGFR monoclonal antibody, especially in those with wild-type *KRAS*.

Finally, Mesteri *et al.* concluded that *BRAF* V600 mutation can be used as a classification criterion for the evaluation of serrated lesions and progression to sessile/serrated adenoma polyps and CRC.

## **Vascular Endothelial Growth Factor (VEGF)**

VEGF is a proangiogenic factor involved in endothelial cell proliferation, migration and vascular permeability (3). Increase in its expression is associated with poor prognosis, low response to preoperative radiotherapy, and greater likelihood of recurrence (43).

Angiogenesis plays a significant role in CRC as neovascularization markers are overexpressed in most of the cancer cells (44). VEGFA is involved in early tumor stages *i.e.* adenoma formation. According to the same study, VEGFA was associated with advanced cancer stage and metastatic disease *via* a angiolymphatic invasion pathway.

Moreover, a relationship between VEGFC and collagen triple helix containing 1 has been noted in terms of prognostic value in rectal cancer (45). VEGFC activates the tyrosine kinase-linked receptor of the VEGFR3 pathway



Table II. *KRAS* and *BRAF* gene mutations baseline characteristics. This table describes the baseline characteristics of *KRAS* and *BRAF* mutations regarding predictive value and common mutation patterns.

Gene	wt/mutant	Baseline characteristics
KRAS	wt	Better response to EGFR inhibitors (1, 3, 9, 28-31)
	Mutant	Most likely at codons 12, 13, 61 (2) In total, 85 KRAS mutation have been identified and occupy around 30-40% of CRC
BRAF	V600E mutation	Predictive value is still under discussion (3, 29) <i>BRAF</i> mutations are related to 10-20% of CRC (Lockhead <i>et al.</i> ) and they are associated with inferior prognosis (13) V600E mutation is associated with sporadic MSI+ tumors (2, 3, 9, 10, 19)

BRAF V600E: v-raf murine sarcoma viral oncogene homolog B V600E mutation; EGFR: epidermal growth factor.

and in-duces lymphangiogenesis. (44) This means that the overexpression of VEGFC could be a valid molecular biomarker for lymphangiogenic metastasis.

### Cyclooxygenase 2 (COX2)-related Pathway

Several prospective studies have shown that COX2 overexpression could be interpreted as an adverse prognostic factor for CRC (46).

There are some studies which support that the use of COX2 inhibitors, *i.e.* aspirin, can be associated with lower risk of CRC, especially in patients who have been diagnosed with CRC in which COX2 is overexpressed (47).

Another study evaluated the correlation between aspirin intake and CRC according to *BRAF* mutation status using questionnaire on aspirin use, and collected data from 1986 to 2006 (48). The study concluded that regular aspirin use was associated with lower risk of wild-type *BRAF* CRC but not of *BRAF*-mutated cancer risk.

Finally, a recent study noted (49) that positivity for COX2 and VEGF was strongly correlated with decreased Disease Free Survival ( $p=0.007$ ), whereas combinations of positivity for RAF kinase inhibitor protein with COX2 negativity, or with negativity for VEGF was strong correlated with improved Disease Free Survival.

### LOH 18q

LOH, especially in 18q, is a well-known potential mechanism for tumorigenesis that has been studied in CRC. LOH is an important mechanism of inactivation of tumor-suppressor genes (3). There have been several studies to interpret the potential prognostic effect of LOH 18q in CRC. LOH in 18q has been inversely associated with MSI (15) There are many tumor-suppressor genes in 18q, *e.g.* *Deleted in Colorectal Carcinoma (DCC)*, *Mothers Against Decapentaplegic Homolog 4 (SMAD4)*, *SMAD2*, and *CDK5* And *ABL1 Enzyme Substrate (CABLES1)* (15) which might interfere with and subsequently explain the prognostic value of LOH 18q.

Nevertheless, in the same study, in 555 non-MSI-high tumor samples with informative 18q LOH data, LOH was present in at least one 18q marker in 362 tumors (65%). There was an association between LOH 18q and obesity (Body mass index  $\geq 30$  kg/m<sup>2</sup>;  $p=0.018$ ), distal colon location ( $p=0.025$ ), low tumor grade ( $p=0.0060$ ), low-level long interspersed nucleotide element-1 methylation (LINE1) methylation ( $p=0.040$ ), wild-type *KRAS* ( $p=0.015$ ), and ohn Cunningham (JC) virus T antigen (JCVT) ( $p=0.0004$ ). The same study concluded that there is no association of LOH 18q with prognosis as yet.

### miRNA – Developing New Promising Biomarkers

miRNAs are short, 18-25 nucleotide non-coding single-stranded RNA sequences which are involved in regulation of gene expression on a post-transcriptional level, through binding to their target protein-encoding mRNA (50). They are believed to play a potentially significant role in the pathogenesis of CRC (51). There are several studies which identify the presence of various miRNAs as predictive or prognostic biomarkers (51-57). In the study of Svoboda *et al.*, miR-215, miR-99a\*, miR-196b, miR-450b-5p and LET-7e were associated with expression of thymidylate synthetase and radioresistance or chemoresistance to its inhibitors (51).

miRNAs are also involved in the regulation of the EGFR signaling pathway (57) and thus they may have essential value in predicting the response to EGFR inhibitors. Another study identified that miR-451 inhibits cell growth through down-regulating the P13K/AKT pathway and thus it potentially has a repressive role (52). Lou *et al.* claimed that miR-625 may have a prognostic and predictive role in CRC as decreased levels are associated with high incidence of metastasis and poor prognosis (53). Yang *et al.* identified miR-29c as having an antitumorigenic role and, preoperatively, decreased levels of miR-29c were associated with CRC relapse. miRNA-126 can have anticarcinogenic properties through the inhibition of neovascularization *via* blocking the 3'-UTR' of VEGF (55).

Karaayvaz *et al.* highlighted that miR-129 could potentially have a tumor-suppressor role and might be a novel target for

Table III. *MicroRNAs involved in CRC mechanisms. This table describes the main microRNAs that are involved in potential CRC pathways. Each might have different mechanisms of action by which it could result in inhibition or acceleration of tumorigenesis.*

miRNA	Mechanism of action
miRNA-451	Inhibits cell growth through the down-regulation of PI3K/AKT pathway (52)
miRNA-625	Decreased levels are associated with high incidence of metastasis (53)
miRNA-126	Anti-cancerous action <i>via</i> blocking the 3-UTR <sup>o</sup> of VEGF and inhibiting neovascularization (55)
miRNA-129	Tumor suppressor role, potential target for therapies (56)
miRNA-133a	Anticancerous properties <i>via</i> increase of the expression of <i>p53</i> and <i>p21</i> gene products (57)
miRNA-224	Can induce tumour expansion <i>via</i> repressing <i>PHLPP1</i> and <i>PHLPP2</i> (58)

PHLPP1: PH domain and leucine rich repeat protein phosphatase 1; PHLPP2: PH domain and leucine rich repeat protein phosphatase 2; VEGF: vascular endothelial growth factor; PI3K/AKT: phosphoinositide-3-kinase/v-akt murine thymoma viral oncogene.

anti-CRC therapies (56). Dong *et al.* claim that miR-133a serves as a functional tumor suppressor in CRC *via* enhancing apoptosis and inhibiting cell proliferation (57). According to the same study, miRNA 133a increased p53 protein and induced *p21* transcription, and can serve as a sensitizer to doxorubicin and oxaliplatin. One potential mechanism of the anticancer properties of miRNA-133a is the fact that it can repress the ring finger and FYVE-like domain containing E3 ubiquitin protein ligase 3'UTR reporter activity and reduce its protein level. Finally, Liao *et al.* noted that miRNA-224 can induce tumor expansion and cell proliferation *via* repressing PH domain and leucine rich repeat protein phosphatase 1 and -2 genes (58). miRNAs seem to be an interesting field of research with promising results (Table III).

### Conclusion

CRC is the second most frequent cancer in Europe and despite progress in the understanding of the molecular background of CRC, effective molecular classification remains a challenge.

Facing these difficulties in the establishment of a key model, researchers have started to try different, more innovative approaches, in which gene signatures dominate. In fact, the idea of gene signatures is very promising (59, 60), but there are still several issues of reproducibility and possible overlap (61, 62). Metabolomics is a new and promising field which tends to utilize simple products from cellular metabolism in order to identify differential and pathognomonic features of cancer (63-66). The main advantage of this approach is the easy access; however, the low specificity for CRC remains a great challenge. Nevertheless, there is much to be looked at in this field, as metabolomics could potentially be a new concept in the investigation and classification of CRC.

Thus, molecular classification of CRC still remains a challenge that researchers have to overcome in the coming years. Creating an effective classification of CRC biomarkers based on their prognostic and predictive value will generate an effective decision-making tool in everyday's clinical practice.

### References

- 1 Ceelen WP: Progress in rectal cancer treatment. *ISRN Gastroenterol* 2012: 648183, 2012.
- 2 Church D, Midgley R and Kerr D: Biomarkers in early-stage colorectal cancer: Ready for prime time? *Dig Dis* 30(2): 27-33, 2012.
- 3 Lee JK and Chan AT: Molecular prognostic and predictive markers in colorectal cancer: Current status. *Curr Colorectal Cancer Rep* 7(2): 136-144, 2011.
- 4 Ogino S and Goel A: Molecular classification and correlates in colorectal cancer. *J Mol Diagn* 10(1): 13-27, 2008.
- 5 Soreide K, Berg M, Skudal BS and Nedreboe BS: Advances in the understanding and treatment of colorectal cancer. *Discov Med* 12(66): 393-404, 2011.
- 6 Bettington M, Walker N, Clouston A, Brown I, Leggett B and Whitehall V: The serrated pathway to colorectal carcinoma: current concepts and challenges. *Histopathology* 62(3): 367-386, 2013.
- 7 Schweiger MR, Hussong M, Röhr C and Lehrach H: Genomics and epigenomics of colorectal cancer. *Wiley Interdiscip Rev Syst Biol Med* 5(2): 205-219, 2013.
- 8 Gyparaki MT, Basdra EK and Papavassiliou AG: DNA methylation biomarkers as diagnostic and prognostic tools in colorectal cancer. *J Mol Med (Berl)* 91(11): 1249-1256, 2013.
- 9 Corso G, Pascale V, Flauti G, Ferrara F, Marrelli D and Roviello F: On-cogenic mutations and microsatellite instability phenotype predict specific anatomical subsite in colorectal cancer patients. *Eur J Hum Genet* 21(12): 1383-1388, 2013.
- 10 Domingo E, Ramamoorthy R, Oukrif D, Rosmarin D, Presz M, Wang H, Pulker H, Lockstone H, Hveem T, Cranston T, Danielsen H, Novelli M, Davidson B, Xu ZZ, Molloy P, Johnstone E, Holmes C, Midgley R, Kerr D, Sieber O and Tomlinson I: Use of multivariate analysis to suggest a new molecular classification of colorectal cancer. *J Pathol* 229(3): 441-448, 2013.
- 11 Markovic S, Antic J, Dimitrijevic I, Zogovic B, Bojic D, Svorcan P, Markovic V and Krivokapic Z: Microsatellite instability affecting the T17 repeats in intron 8 of HSP110, as well as five mononucleotide repeats in patients with colorectal carcinoma. *Biomark Med* 7(4): 613-621, 2013.
- 12 Budinska E, Popovici V, Tejpar S, D'Ario G, Lapique N, Sikora KO, Di Narzo AF, Yan P, Hodgson JG, Weinrich S, Bosman F, Roth A and Delorenzi M: Gene expression patterns unveil a new level of molecular heterogeneity in colorectal cancer. *J Pathol* 231(1): 63-76, 2013.

- 13 Lochhead P, Kuchiba A, Imamura Y, Liao X, Yamauchi M, Nishihara R, Qian ZR, Morikawa T, Shen J, Meyerhardt JA, Fuchs CS and Ogino S: Microsatellite instability and BRAF mutation testing in colorectal cancer prognostication. *J Natl Cancer Inst* 105(15): 1151-1156, 2013.
- 14 Marisa L, de Reyniès A, Duval A, Selves J, Gaub MP, Vescovo L, Etienne-Grimaldi MC, Schiappa R, Guenot D, Ayadi M, Kirzin S, Chazal M, Fléjou JF, Benchimol D, Berger A, Lagarde A, Pencreach E, Piard F, Elias D, Parc Y, Olschwang S, Milano G, Laurent-Puig P and B expression classification of colon cancer into molecular sub-types: characterization, validation, and prognostic value. *PLoS Med* 10(5): e1001453, 2013.
- 15 Ogino S1 Noshō K, Irahara N, Shima K, Baba Y, Kirkner GJ, Meyerhardt JA and Fuchs CS: Prognostic significance and molecular associations of 18q loss of heterozygosity: a cohort study of microsatellite stable colorectal cancers. *J Clin Oncol* 27(27): 4591-4598, 2009.
- 16 Thompson BA1, Goldgar DE, Paterson C, Clendenning M, Walters R, Arnold S, Parsons MT, Michael D W, Gallinger S, Haile RW, Hopper JL, Jenkins MA, Lemarchand L, Lindor NM, Newcomb PA, Thibodeau SN; Colon Cancer Family Registry, Young JP, Buchanan DD, Tavtigian SV and Spurdle AB: A multifactorial likelihood model for MMR gene variant classification incorporating probabilities based on sequence bioinformatics and tumor characteristics: a report from the Colon Cancer Family Registry. *Hum Mutat* 34(1): 200-209, 2013.
- 17 Losso GM, Moraes Rda S, Gentili AC and Messias-Reason IT: Micro-satellite instability – MSI markers (BAT26, BAT25, D2S123, D5S346, D17S250) in rectal cancer. *Arq Bras Cir Dig* 25(4): 240-244, 2012.
- 18 Jass JR: Classification of colorectal cancer based on correlation of clinical, morphological and molecular features. *Histopathology* 50(1): 113-130, 2007.
- 19 Simons CC, Hughes LA, Smits KM, Khalid-de Bakker CA, de Bruïne AP, Carvalho B, Meijer GA, Schouten LJ, van den Brandt PA, Weijenberg MP and van Engeland M: A novel classification of colorectal tumors based on microsatellite instability, the CpG island methylator phenotype and chromosomal instability: implications for prognosis. *Ann Oncol* 24(8): 2048-2056, 2013.
- 20 Haglund HR, Berg M, Jolma IW, Carlsen A and Søreide K: Molecular Pathways and Cellular Metabolism in Colorectal Cancer. *Dig Surg* 30(1): 12-25, 2013.
- 21 Tomlinson I, Halford S, Aaltonen L, Hawkins N and Ward R: Does MSI-low exist? *J Pathol* 197(1): 6-13, 2002.
- 22 Gaiser T, Meinhardt S, Hirsch D, Killian JK, Gaedcke J, Jo P, Ponsa I, Miró R, Rüschoff J, Seitz G, Hu Y, Camps J and Ried T: Molecular patterns in the evolution of serrated lesion of the colorectum. *Int J Cancer* 132(8): 1800-1810, 2013.
- 23 Migliore L, Migheli F, Spisni R and Coppedè F: Genetics, cytogenetics, and epigenetics of colorectal cancer. *J Biomed Biotechnol* 2011: 792362, 2011.
- 24 Gerling M, Glauben R, Habermann JK, Kühl AA, Loddenkemper C, Lehr HA, Zeitz M and Siegmund B: Characterization of chromosomal instability in murine colitis-associated colorectal cancer. *PLoS One* 6(7): e22114, 2011.
- 25 Hershkovitz D and Ben-Izhak O: Molecular pathology - aspects in KRAS mutation analysis in colon carcinoma. *Harefuah* 152(6): 356-360 and 367, 2013.
- 26 Phipps AI, Buchanan DD, Makar KW, Win AK, Baron JA, Lindor NM, Potter JD and Newcomb PA: KRAS-mutation status in relation to colo-rectal cancer survival: the joint impact of correlated tumour markers. *Br J Cancer* 108(8): 1757-1764, 2013.
- 27 Marchoudi N, Amrani Hassani Joutei H, Jouali F, Fekkek J and Rhaissi H: Distribution of KRAS and BRAF mutations in Moroccan patients with advanced colorectal cancer. *Pathol Biol (Paris)* 61(6): 273-276, 2013.
- 28 Grade M, Wolff HA, Gaedcke J and Ghadimi BM: The molecular basis of chemoradiosensitivity in rectal cancer: implications for personalized therapies. *Langenbecks Arch Surg* 397(4): 543-555, 2012.
- 29 Rizzo S, Bronte G, Fanale D, Corsini L, Silvestris N, Santini D, Gulotta G, Bazan V, Gebbia N, Fulfaro F and Russo A: Prognostic vs. predictive molecular biomarkers in colorectal cancer: is KRAS and BRAF wild type status required for anti-EGFR therapy? *Cancer Treat Rev* 36(3): S56-61, 2010.
- 30 Siena S, Sartore-Bianchi A, Di Nicolantonio F, Balfour J and Bardelli A: Biomarkers predicting clinical outcome of epidermal growth factor re-ceptor-targeted therapy in metastatic colorectal cancer. *J Natl Cancer Inst* 101(19): 1308-1324, 2009.
- 31 Duffy MJ, Lamerz R, Haglund C, Nicolini A, Kalousová M, Holubec L and Sturgeon C: Tumor markers in colorectal cancer, gastric cancer and gastrointestinal stromal cancers: European group on tumor markers 2014 guidelines update. *Int J Cancer*. 2013 doi: 10.1002/ijc.28384. [Epub ahead of print]
- 32 Di Bartolomeo M, Pietrantonio F, Perrone F, Dotti KF, Lampis A, Bertan C, Beretta E, Rimassa L, Carbone C, Biondani P, Passalacqua R, Pilotti S and Bajetta E; on behalf of Italian Trials in Medical Oncology (ITMO) Group: Lack of KRAS, NRAS, BRAF and TP53 mutations improves outcome of elderly metastatic colorectal cancer patients treated with cetuximab, oxaliplatin and UFT. *Target Oncol*. 2013 [Epub ahead of print].
- 33 Selcukbiricik F, Erdamar S, Ozkurt CU, Molinas Mandel N, Demirelli F, Ozguroglu M, Tural D, Buyukunal E and Serdengeci S: The role of K-RAS and B-RAF mutations as biomarkers in metastatic colorectal cancer. *J BUON* 18(1): 116-123, 2013.
- 34 Bruera G, Cannita K, Di Giacomo D, Lamy A, Frébourg T, Sabourin JC, Tosi M, Alesse E, Ficorella C and Ricevuto E: Worse prognosis of KRAS c.35 G>A mutant metastatic colorectal cancer (MCRC) patients treated with intensive triplet chemotherapy plus bevacizumab (FIR-B/FOX). *BMC Med* 11: 59, 2013.
- 35 Loupakis F, Cremolini C, Fioravanti A, Orlandi P, Salvatore L, Masi G, Schirripa M, Di Desidero T, Antoniotti C, Canu B, Faviana P, Sensi E, Lupi C, Fontanini G, Basolo F, Di Paolo A, Danesi R, Falcone A and Bocci G: EGFR ligands as pharmacodynamic biomarkers in metastatic colorectal cancer patients treated with cetuximab and irinotecan. *Target Oncol* 2013 [Epub ahead of print].
- 36 Tian S, Simon I, Moreno V, Roepman P, Tabernero J, Snel M, van't Veer L, Salazar R, Bernards R and Capella G: A combined oncogenic pathway signature of BRAF, KRAS and PI3KCA mutation improves colorectal cancer classification and cetuximab treatment prediction. *Gut* 62(4): 540-549, 2013.
- 37 Saridaki Z, Tzardi M, Papadaki C, Sfakianaki M, Pega F, Kalikaki A, Tsakalaki E, Trypaki M, Messaritakis I, Stathopoulos E, Mavroudis D, Georgoulas V and Souglakos J: Impact of KRAS, BRAF, PIK3CA mutations, PTEN, AREG, EREG expression and skin rash in ≥ 2 line cetuximab-based therapy of colorectal cancer patients. *PLoS One* 6(1): e15980, 2011.

- 38 Lièvre A, Rouleau E, Buecher B and Mitry E: Clinical significance of BRAF mutations in colorectal cancer. *Bull Cancer* 97(12): 1441-1452, 2010.
- 39 Adackapara CA, Sholl LM, Barletta JA and Hornick JL: Immunohisto-chemistry using the BRAF V600E mutation-specific monoclonal anti-body VE1 is not a useful surrogate for genotyping in colorectal adeno-carcinoma. *Histopathology* 63(2): 187-193, 2013.
- 40 Eklöf V, Wikberg ML, Edin S, Dahlin AM, Jonsson BA, Öberg Å, Rutegård J and Palmqvist R: The prognostic role of KRAS, BRAF, PIK3CA and PTEN in colorectal cancer. *Br J Cancer* 108(10): 2153-2163, 2013.
- 41 Yokota T, Ura T, Shibata N, Takahari D, Shitara K, Nomura M, Kondo C, Mizota A, Utsunomiya S, Muro K and Yatabe Y: BRAF mutation is a powerful prognostic factor in advanced and recurrent colorectal cancer. *Br J Cancer* 104(5): 856-862, 2011.
- 42 Yuan ZX, Wang XY, Qin QY, Chen DF, Zhong QH, Wang L and Wang JP: The prognostic role of BRAF mutation in metastatic colorectal cancer receiving anti-EGFR monoclonal antibodies: a meta-analysis. *PLoS One* 8(6): e65995, 2013.
- 43 Zlobec I, Vuong T, Compton CC, Lugli A, Michel RP, Hayashi S and Jass JR: Combined analysis of VEGF and EGFR predicts complete tumour response in rectal cancer treated with preoperative radiotherapy. *Br J Cancer* 98(2): 450-456, 2013.
- 44 Martins SF, Garcia EA, Luz MA, Pardo F, Rodrigues M and Filho AL: Clinicopathological correlation and prognostic significance of VEGF-A, VEGF-C, VEGFR-2 and VEGFR-3 expression in colorectal cancer. *Cancer Genomics Proteomics* 10(2): 55-67, 2013.
- 45 Yin WH, Fan HZ, Sheng JW, Xia HM, Wu YW and Xie P: Effect of vascular endothelial growth factor C and collagen triple helix repeat con-taining 1 expression on prognosis of rectal carcinoma patients. *Zhonghua Wei Chang Wai Ke Za Zhi* 16(7): 673-675, 2013.
- 46 Soumaoro LT, Uetake H, Higuchi T, Takagi Y, Enomoto M and Sugi-hara K: Cyclooxygenase-2 expression: a significant prognostic indica-tor for patients with colorectal cancer. *Clin Cancer Res* 10(24): 8465-8471, 2004.
- 47 Chan AT, Ogino S and Fuchs CS: Aspirin use and survival after diag-nosis of colorectal cancer. *JAMA* 302(6): 649-658, 2009.
- 48 Nishihara R, Lochhead P, Kuchiba A, Jung S, Yamauchi M, Liao X, Imamura Y, Qian ZR, Morikawa T, Wang M, Spiegelman D, Cho E, Giovannucci E, Fuchs CS, Chan AT and Ogino S: Aspirin use and risk of colorectal cancer according to BRAF mutation status. *JAMA* 309(24): 2563-2571, 2013.
- 49 Kim JW, Kim YB, Choi JJ, Koom WS, Kim H, Kim NK, Ahn JB, Lee I, Cho JH and Keum KC: Molecular markers predict distant metastases after adjuvant chemoradiation for rectal cancer. *Int J Radiat Oncol Biol Phys* 84(5): e577-584, 2012.
- 50 Mlcochova J, Faltejskova P, Nemecek R, Svoboda M and Slaby O: Mi-croRNAs targeting EGFR signalling pathway in colorectal cancer. *J Cancer Res Clin Oncol* 139(10): 1615-1624, 2013.
- 51 Svoboda M, Sana J, Fabian P, Kocakova I, Gombosova J, Nekvindova J, Radova L, Vyzula R and Slaby O: MicroRNA expression profile as-sociated with response to neoadjuvant chemoradiotherapy in locally advanced rectal cancer patients. *Radiat Oncol* 7: 195, 2012.
- 52 Li HY, Zhang Y, Cai JH and Bian HL: MicroRNA-451 inhibits growth of human colorectal carcinoma cells *via* down-regulation of PI3K/AKT pathway. *Asian Pac J Cancer Prev* 14(6): 3631-3634, 2013.
- 53 Lou X, Qi X, Zhang Y, Long H and Yang J: Decreased expression of microRNA-625 is associated with tumor metastasis and poor prognosis in patients with colorectal cancer. *J Surg Oncol* 108(4): 230-235, 2013.
- 54 Yang IP, Tsai HL, Huang CW, Huang MY, Hou MF, Juo SH and Wang JY: The functional significance of microRNA-29c in patients with colo-rectal cancer: a potential circulating biomarker for predicting early re-lapse. *PLoS One* 8(6): e66842, 2013.
- 55 Zhang Y, Wang X, Xu B, Wang B, Wang Z, Liang Y, Zhou J, Hu J and Jiang B: Epigenetic silencing of miR-126 contributes to tumor invasion and angiogenesis in colorectal cancer. *Oncol Rep* 30(4): 1976-1984, 2013.
- 56 Karaayvaz M, Zhai H and Ju J: miR-129 promotes apoptosis and en-hances chemosensitivity to 5-fluorouracil in colorectal cancer. *Cell Death Dis* 4: e659, 2013.
- 57 Dong Y, Zhao J, Wu CW, Zhang L, Liu X, Kang W, Leung WW, Zhang N, Chan FK, Sung JJ, Ng SS and Yu J: Tumor suppressor functions of miR-133a in colorectal cancer. *Mol Cancer Res* 11(9): 1051-1060, 2013.
- 58 Liao WT, Li TT, Wang ZG, Wang SY, He MR, Ye YP, Qi L, Cui YM, Wu P, Jiao HL, Zhang C, Xie YJ, Wang JX and Ding YQ: microRNA-224 promotes cell proliferation and tumor growth in human colorectal can-cer by repressing PHLPP1 and PHLPP2. *Clin Cancer Res* 19(17): 4662-4672, 2013.
- 59 Kalady MF, DeJulius K, Church JM, Lavery IC, Fazio VW and Ishwaran H: Gene signature is associated with early-stage rectal cancer recur-rence. *J Am Coll Surg* 211(2): 187-195, 2010.
- 60 Wang Y, Jatke T, Zhang Y, Mutch MG, Talantov D, Jiang J, McLeod HL and Atkins D: Gene expression profiles and molecular markers to predict recurrence of Dukes' B colon cancer. *J Clin Oncol* 22(9): 1564-1571, 2004.
- 61 Sanz-Pamplona R, Berenguer A, Cordero D, Riccadonna S, Solé X, Crous-Bou M, Guinó E, Sanjuan X, Biondo S, Soriano A, Jurman G, Capella G, Furlanello C and Moreno V: Clinical value of prognosis gene expression signatures in colorectal cancer: a systematic review. *PLoS One* 7(11): e48877, 2012.
- 62 Akiyoshi T, Kobunai T and Watanabe T: Predicting the response to preoperative radiation or chemoradiation by a microarray analysis of the gene expression profiles in rectal cancer. *Surg Today* 42(8): 713-719, 2012.
- 63 Ma Y, Zhang P, Wang F, Liu W, Yang J and Qin H: An integrated pro-teomics and metabolomics approach for defining oncofetal biomarkers in the colorectal cancer. *Ann Surg* 255(4): 720-730, 2012.
- 64 Warburg O: On the origin of cancer cells. *Science* 123(3191): 309-314, 1956.
- 65 DeBerardinis RJ, Mancuso A, Daikhin E, Nissim I, Yudkoff M, Wehrli S and Thompson CB: Beyond aerobic glycolysis: transformed cells can engage in glutamine metabolism that exceeds the requirement for pro-tein and nucleotide synthesis. *Proc Natl Acad Sci USA* 104(49): 19345-19350, 2007.
- 66 Vermeersch KA and Styczynski MP: Applications of meta-bolomics in cancer research. *J Carcinog* 12: 9, 2013.

Received January 22 2014  
 Revised March 5, 2014  
 Accepted March 7, 2014