

Effects of Antioxidants and Pro-oxidants on Cytotoxicity of Dihydroartemisinin to Molt-4 Human Leukemia Cells

THOMAS GERHARDT, ROSS JONES, JUNGSOO PARK, RUI LU,
HO WING CHAN, QINGXI FANG, NARENDRA SINGH and HENRY LAI

Department of Bioengineering, University of Washington, Seattle, WA, U.S.A.

Abstract. *Background: The objective of the present study was to investigate how oxidative status influences the effectiveness of cytotoxicity of artemisinin towards cancer cells. It is hypothesized that antioxidants would reduce, whereas pro-oxidants would enhance, cytotoxicity. Materials and Methods: Molt-4 human leukemia cells were incubated with vitamins C, E, D₃, dexamethasone, or hydrogen peroxide alone or in combination with dihydroartemisinin (DHA). Concentrations of these compounds studied were similar to those achievable by oral administration. Viable cell counts were performed before (0 h) and at, 24 and 48 h after treatment. Results: Vitamin C, vitamin D₃, dexamethasone, and H₂O₂ caused significant Molt-4 cell death. Vitamin E caused an increase in Molt-4 cell growth. Vitamin C and vitamin D₃ significantly interacted with DHA at the 48-h time point and with H₂O₂ at both 24-h and 48-h time points. Conclusion: Cellular oxidative status could alter the potency of artemisinin in killing cancer cells.*

Environmental factors, such as nutrition and nutritional supplementation, can explain some of the variability seen among treatment outcomes of cancer patients (1). To explore this possibility, we tested the effects of several nutritional supplements commonly taken by cancer patients on an experimental cancer drug artemisinin, which we have been investigating as an alternative to current chemotherapeutics (2). Artemisinin, a compound isolated from the plant *Artemisia annua* L (sweet wormwood), and its derivatives are currently the most effective anti-malarials (3). Artemisinin compounds have also been shown to be potent and selective anticancer

Correspondence to: Dr. Narendra Singh, Department of Bioengineering, Box 355061, University of Washington, Seattle, WA 98195-5061, U.S.A. Tel/Fax: +1 2066852060, e-mail: narendra@u.washington.edu

Key Words: Artemisinin, sodium ascorbate, vitamin D₃, Trolox, dexamethasone, hydrogen peroxide, Molt-4 cells.

agents both *in vitro* and *in vivo* (2). Artemisinin's anticancer mechanism has been hypothesized to arise from the interaction of its endoperoxide moiety with free ferrous iron, which is more abundant in cancer cells, generating cytotoxic reactive oxygen species (ROS) (4, 5). There are several case reports of cancer treatment using artemisinin (6-8).

In the present study, we investigated whether various nutritional supplements interact with the more hydrophilic artemisinin analog dihydroartemisinin (DHA) in killing cancer cells. Since artemisinin works by overloading cancer cells with ROS, we selected molecules that are known to affect the oxidative status of cells. For their antioxidant properties, we studied sodium ascorbate (vitamin C) and Trolox (vitamin E). For their pro-oxidant properties, in certain cancer cells, we studied vitamin D₃, dexamethasone (dexamethasone 21-acetate), and hydrogen peroxide (H₂O₂). To evaluate the interactions of artemisinin with these agents, we used the commonly studied, rapidly growing and easily cultured Molt-4 human lymphoblastic leukemia cells.

Materials and Methods

Chemicals. All chemicals were purchased from Sigma Aldrich (St. Louis, MO, USA) unless mentioned otherwise.

Molt-4 cell culture. Molt-4 cells were purchased from American Type Culture Collection (ATCC, Manassas, VA, USA). They were cultured in RPMI-1640 media (Life Technologies, Grand Island, NY, USA) with 10% fetal bovine serum (ATCC) at 37°C with 5% CO₂ and 100% humidity to reach a density of 6×10⁵ cells/ml before diluting to a density of 6×10⁴ cells/ml approximately 24 h prior to a treatment. Twenty-four hours of pre-incubation allowed the cells and media to be conditioned prior to exposure to supplements and experiments to begin at a density of approximately 1×10⁵ cells/ml. At this time, cells were in the log phase. Viable cells were counted before drug treatment (0 h) and at 24 h and 48 h after treatment using a hemocytometer. To assess cell viability, we used trypan blue exclusion and identified dead cells based upon morphological criteria: chromatin condensation, formation of apoptotic bodies, shrinkage of the cytoplasm, and blebbing of the plasma membrane with irregular outline (9, 10). Only healthy cells were counted.

Table I. Results of two-way ANOVA showing significant differences of dihydroartemisinin (DHA), supplements (ascorbate, Vitamin D₃, dexamethasone, Trolox and hydrogen peroxide) and interaction (DHA+supplements) effects at the 24-h and 48-h time points. Results showing significance are in bold.

Supplement	Treatment	Time (h)	n	p (DHA)	p (supplement)	p (Interaction)
Ascorbate	256 μM	24	3	0.0008	0.0053	0.4762
		48	3	<0.0001	0.0001	0.0071
Vitamin D ₃	75 nM	24	3	0.0009	0.8118	0.9924
		48	3	<0.0001	0.0072	0.0096
Trolox	45.8 μM	24	3	0.0021	0.0175	0.5329
		48	3	<0.0001	0.6109	0.7447
Dexamethasone	10 μM	24	4	<0.0001	0.0448	0.6221
		48	4	<0.0001	0.0022	0.5054
Hydrogen Peroxide	8.82 μM	24	3	<0.0001	<0.0001	0.0331
		48	3	<0.0001	<0.0001	<0.0001

Cell treatment. There were four treatment groups: control (no-drug treatment), Dihydroartemisinin (DHA) alone, supplement alone, and DHA+supplement. DHA (Holley Pharmaceuticals, Chongqing, China) was used at a concentration of 12.4 μM. This concentration was used because it is similar to the maximum blood concentration in rats after an oral dosage of 10 mg/kg (11). It was also the concentration used in our previous studies. The supplement samples were added with the following concentrations: ascorbate 256 μM, vitamin D₃ 75 nM, Trolox 45.8 μM, dexamethasone 21-acetate 10 μM, and H₂O₂ 8.82 μM, according to standard achievable *in vivo* concentrations of these compounds. Dimethyl sulfoxide (DMSO) was the solvent for DHA, dexamethasone, Trolox, and vitamin D₃. All other compounds were dissolved in RPMI-1640 media. All groups were controlled at 1% DMSO by adding the appropriate amount of DMSO directly to the suspension culture at the experimental initiation. Each experiment was performed at least three times.

Data analysis. Data were expressed as the ratio of cell counts at the 24-h or 48-h time point relative to time zero. The GraphPad Prism 6.03 software (La Jolla, CA, USA) was used for statistical analysis. A two-way ANOVA was performed to test for significances of main effects of DHA and supplements, as well as their interaction effect. Pair-wise comparisons were analyzed using the Newman Keul's test. A *p*<0.05 was considered statistically significant.

Results

Data of effects of DHA and different supplements on Molt-4 cells are presented in Figure 1. Results of two-way ANOVA showing significances of main and interaction effects are presented in Table I.

In all cases, DHA-alone significantly decreased cell counts relative to untreated cells at both 24-h and 48-h time points. All supplements studied also affected cell counts. Compared to control, ascorbate and H₂O₂ both caused a significant decrease in cell count both at 24-h (*p*<0.05 and *p*<0.0001 for ascorbate and H₂O₂, respectively) and 48-h (*p*<0.0001 for both ascorbate and H₂O₂) time points. Dexamethasone and vitamin D₃ caused a significant decrease in cell count only at the 48-h time point (*p*<0.01 for both supplements).

A significant increase in cell count was observed at the 24-h time point for Trolox (*p*<0.05). Thus, all supplements, except for Trolox, caused a decrease in cell counts.

Hydrogen peroxide in combination with DHA caused significantly greater cell death than DHA-alone at both time points (*p*<0.01 and *p*<0.05 for 24-h and 48-h time points, respectively). Also, the effect of dexamethasone combined with DHA was found to be significantly greater than DHA alone at the 48-h time point (*p*<0.05). However, cell counts of ascorbate, vitamin D₃ or Trolox in combination with DHA are not significantly different from DHA alone at both 24-h and 48-h time points. This is reflected in the results of the two-way ANOVA (Table I) indicating that H₂O₂ significantly interacted with DHA at both time points, whereas ascorbate and Vitamin D₃ were found to interact at the 48-h time point. No significant interaction effect was observed between DHA and Trolox or dexamethasone.

Discussion

The effectiveness of artemisinin may be due to its free radical generation capabilities in the presence of Fe²⁺ causing widespread effects on cancer cells (12, 13). If patients are undergoing cancer therapies with drugs like artemisinin, which rely on oxidative stress, then it is important to know how supplementation with oxidative-state modulating compounds will affect the therapy. It may be possible to improve therapeutic outcomes by compounding the oxidizing effect of artemisinin and certain supplements, pushing cancer cells toward oxidative overload (14).

Vitamin C, a common supplement, has been shown to act as both a ROS generator in the presence of iron and copper (15) and as an antioxidant (16). The normal blood concentration of vitamin C is approximately 64 μM in an adult man (17). We hypothesize that ascorbate acts more like a pro-oxidant in cancer cells due to their higher cytoplasmic iron levels. From our results, we can postulate that ROS




Figure 1. Effects of dihydroartemisinin (DHA), supplements, and DHA+supplements on Molt-4 cells at 24-h and 48-h time points. Error bar denotes standard deviation. *: $p < 0.05$, **: $p < 0.01$, ****: $p < 0.0001$ comparing supplement with control. †: $p < 0.05$, ††: $p < 0.01$ comparing DHA+ supplement with DHA.

generation is causing cell death independently and in combination with DHA. Variation in cancer therapy outcomes with drugs targeting oxidative stress may be caused, in part, by vitamin C levels, since it is a commonly used supplement.

However, one must be careful in taking vitamin C in combination with DHA. Ascorbate can convert ferric iron into ferrous iron (18), the active form that reacts with artemisinin, generating short lived free radicals. If this happens in the stomach of a person who is consuming artemisinin along with ascorbate, ascorbate will convert ferric iron in foods to the ferrous form, which may react with artemisinin locally, making the therapy less effective. To avoid this issue, ascorbate may be administered intravenously or taken orally at a different time of day as artemisinin.

Vitamin D₃ has been shown to act as a pro-oxidant and an antioxidant in malignant and non-malignant human prostate epithelial cells, respectively (19). Vitamin D₃ concentrations range between 75 and 190 nM in the blood (20), varying with an individual's age, sunlight exposure, race and other supplement intake (21). Our results indicate that vitamin D₃ reduces leukemia cell growth independently but has no significant interaction effect with DHA. The reason of the lack of an interaction effect is unknown. Perhaps, vitamin D₃ becomes an antioxidant in cancer cells at longer time points. It may be of interest to study the interaction of vitamin D₃ levels with artemisinin *in vivo*, as vitamin D₃ is involved in regulation of calcium and phosphorus intake (22), immune function (23) and, thus, may be involved in more complex interactions with DHA in cancer cells.

The normal blood concentration of vitamin E is 22.9 μM, with a range of 12 to 59 μM (25). Our results with the water-soluble vitamin E analog, Trolox, alone show an increase in leukemia cell growth at the 24-h time point, with no significant interaction with DHA. Vitamin E has been shown to decrease the anti-angiogenic effects of artemisinin (26); thus, it is pertinent to study how it interacts with DHA *in vivo*, where angiogenesis is a major factor in tumor growth.

Dexamethasone is shown to induce oxidative stress-mediated apoptosis in lymphoid tissue (27). Our results demonstrate that dexamethasone reduces leukemia cell growth. There was no significant interaction with DHA. Due to the anti-inflammatory nature of dexamethasone, it will be important to further study its interaction with DHA *in vivo*, where inflammation is a key factor in tumor modulation by the immune system.

H₂O₂ is a known signaling molecule and strong oxidizing agent. Normal human plasma concentration of H₂O₂ is approximately 11.1 μM (31) and can be increased with exercise (32). Our H₂O₂ results indicate that H₂O₂ kills leukemia cells alone. Furthermore, the interaction between H₂O₂ and DHA was found to be additive, which may be due to the similar methods of apoptosis mediation by ROS generation employed by both compounds.

We have explored the interaction effects of several nutritional supplements and natural compounds with antioxidant and pro-oxidant properties with DHA on Molt-4 cells. Some of these compounds could increase or decrease the anticancer effectiveness of artemisinin *in vitro*. We do not yet know the therapeutic values of our *in vitro* findings; however, implications of this work could pave the way for setting new treatment regimens for patients undergoing artemisinin therapy for cancer. Our goal is to understand variability between patients and, hopefully, use this knowledge to reduce the easily achievable dosage of artemisinin required for successful cancer treatment. Our strategy presents an attractive option for patients who are looking to optimize their treatments through nutritional modulation. Future *in vivo* research is needed to confirm our findings in animals and humans and to study more complex systemic interactions with the supplement molecules.

Conflicts of Interest

The Authors declare no conflicts of interest with respect to this study.

Funding

Funding for this research was provided in part by the Dean of the College of Engineering and Associate Vice Provost for Research of the University of Washington and the Washington State Life Science Discovery Fund (LSDF).

References

- 1 Lagiou P, Trichopoulou A and Trichopoulos D: Nutritional epidemiology of cancer: accomplishments and prospects. *Proc Nutr Soc* 61: 217-222, 2002.
- 2 Lai HC, Singh NP and Sasaki T: Development of artemisinin compounds for cancer treatment. *Invest New Drugs* 31: 230-246, 2013.
- 3 White NJ: Assessment of the pharmacodynamic properties of antimalarial drugs *in vivo*. *Antimicrob Agents Chemother* 41: 1413-1422, 1997.
- 4 Lai HC and Singh NP: Selective cancer cell cytotoxicity from exposure to dihydroartemisinin and holotransferrin. *Cancer Lett* 91: 41-46, 1995.
- 5 Singh NP and Lai HC: Selective toxicity of dihydroartemisinin and holotransferrin toward human breast cancer cells. *Life Sci* 70: 49-56, 2001.
- 6 Singh NP and Verma KB: Case report of a laryngeal squamous cell carcinoma treated with artesunate. *Arch Onc* 10: 279-280, 2002.
- 7 Singh NP and Panwar VK: Case report of a pituitary macroadenoma treated with artemether. *Integr Cancer Ther* 5: 391-394, 2006.
- 8 Berger TG, Dieckmann D, Efferth T, Schultz ES, Funk JO, Baur A and Schuler G: Artesunate in the treatment of metastatic uveal melanoma--first experiences. *Oncol Rep* 14: 1599-1603, 2005.
- 9 Searle J, Kerr JFR and Bishop CJ: Necrosis and apoptosis: distinct modes of cell death with fundamentally different significance. *Pathol Ann* 17: 229-259, 1982.

- 10 Kerr JFR, Wyllie AH and Currie AR: Apoptosis: a Basic biological phenomenon with wide-ranging applications in tissue kinetics. *Br J Cancer* 26: 239-257, 1972.
- 11 Li QG, Peggins JO, Fleckenstein LL, Masonic K, Heiffer MH and Brewer TG: The pharmacokinetics and bioavailability of dihydroartemisinin, artether, artemether, artesunic acid and arteminic acid in rats. *J Pharm Pharmacol* 50: 173-182, 1998.
- 12 O'Neill PM, Barton VE and Ward SA: The molecular mechanism of action of artemisinin--the debate continues. *Molecules* 15: 1705-1721, 2010.
- 13 Chan HW, Singh NP and Lai HC. Cytotoxicity of dihydroartemisinin toward Molt-4 cells attenuated by N-tert-butyl-alpha-phenylnitron and deferoxamine. *Anticancer Res* 33: 4389-4393, 2013.
- 14 Schumacker PT: Reactive oxygen species in cancer cells: live by the sword, die by the sword. *Cancer Cell* 10: 175-176, 2006.
- 15 Buettner GR and Jurkiewicz BA: Catalytic metals, ascorbate and free radicals: combinations to avoid. *Radiat Res* 145: 532-541, 1996.
- 16 Gao P, Zhang H, Dinavahi R, Li F, Xiang Y, Raman V, Bhujwalla ZM, Felsher DW, Cheng L, Pevsner J, Lee LA, Semenza GL and Dang CV: HIF-dependent antitumorigenic effect of antioxidants *in vivo*. *Cancer Cell* 12: 230-238, 2007.
- 17 Paszkowski T and Clarke R: The Graafian follicle is a site of L-ascorbate accumulation. *J Assist Reprod Genet* 16: 41-45, 1999.
- 18 Sharp P and Srail SK: Molecular mechanisms involved in intestinal iron absorption. *World J Gastroenterol* 13: 4716-4724, 2007.
- 19 Bao BY, Ting HJ, Hsu JW and Lee YF: Protective role of 1 alpha, 25-dihydroxyvitamin D₃ against oxidative stress in non-malignant human prostate epithelial cells. *Int J Cancer* 122: 2699-2706.
- 20 National Library of Medicine. 25-hydroxy vitamin D test. Available at: <http://www.nlm.nih.gov/medlineplus/ency/article/003569.htm>. Accessibility verified October 29, 2013.
- 21 Weng FL, Shults J, Leonard MB, Stallings VA and Zemel BS: Risk factors for low serum 25-hydroxyvitamin D concentrations in otherwise healthy children and adolescents. *Am J Clin Nutr* 86: 150-158, 2007.
- 22 Holick MF: Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* 80: 1678S-1688S, 2004.
- 23 Bikle DD: Vitamin D regulation of immune function. *Vitam Horm* 86: 1-21, 2011.
- 24 Garcion E, Wion-Barbot N, Montero-Menei CN, Berger F and Wion D: New clues about vitamin D functions in the nervous system. *Trends Endocrinol Metab* 13: 100-105, 2002
- 25 Goraca A: Assessment of total antioxidant capacity in human plasma. *Folia Medica* 46: 16-21, 2004.
- 26 Wartenberg M, Wolf S, Budde P, Grünheck F, Acker H, Hescheler J, Wartenberg G and Sauer H: The antimalaria agent artemisinin exerts antiangiogenic effects in mouse embryonic stem cell-derived embryoid bodies. *Lab Invest* 83: 1647-1655, 2003.
- 27 Baker AF, Briehl MM, Dorr R and Powis G: Decreased antioxidant defense and increased oxidant stress during dexamethasone-induced apoptosis: bcl-2 prevents the loss of antioxidant enzyme activity. *Cell Death Differ* 3: 207-213, 1996.
- 28 Enoiu M, Aberkane H, Salazar JF, Leroy P, Groffen J, Siest G and Wellman M: Evidence for the pro-oxidant effect of gamma-glutamyltranspeptidase-related enzyme. *Free Radic Biol Med* 29: 825-833, 2000.
- 29 Meshnick SR, Thomas A, Ran A, Xy CM and Pan HZ: Artemisinin (qinghaosu): the role of intracellular hemin in its mechanism of antimalarial action. *Mol Biochem Parasitol* 49: 181-189, 1991.
- 30 Michaelis M, Kleinschmidt MC, Barth S, Rothweiler F, Geiler J, Breitling R, Mayer B, Deubzer H, Witt O, Kreuter J, Doerr HW, Cinatl J, Cinatl J Jr. : Anti-cancer effects of artesunate in a panel of chemoresistant neuroblastoma cell lines. *Biochem Pharma* 79: 130-136, 2010.
- 31 Russell IJ, Xiao Y, Haynes WL and Michalek JE: Serum hydrogen peroxide levels are elevated in fibromyalgia syndrome [abstract]. *Arthritis Rheum* 60: Suppl 10:92, 2009.
- 32 Deskur E, Przywarska I, Szcześniak L, Rychlewski T, Wilk M and Wysocki H: Exercise-induced increase in hydrogen peroxide plasma levels is diminished by endurance training after myocardial infarction. *Int J Cardiol* 67: 219-224, 1998.

Received December 15, 2014

Revised January 14, 2015

Accepted January 16, 2015