Biomarkers of Oxidative Stress Associated with the Risk of Potentially Malignant Oral Disorders

THOMAS SENGHORE^{1,2}, YU-FENG LI³, FUNG-CHANG SUNG⁴, MING-HSUI TSAI⁵,
CHUN-HUNG HUA⁵, CHIU-SHONG LIU⁶, RU-JIUN HUANG¹ and CHIH-CHING YEH^{1,7}

¹*School of Public Health, College of Public Health, Taipei Medical University, Taipei, Taiwan, R.O.C.;*

²*Department of Nursing, School of Medicine and Allied Health Sciences,
University of The Gambia, Banjul, Gambia;*

³*Graduate Institute of Biostatistics, Departments of* ⁴*Health Services Administration and
Departments of* ⁵*Otolaryngology, and* ⁶*Family Medicine, China Medical
University Hospital, Taichung, Taiwan, R.O.C.;*

⁷*Public Health, College of Public Health, China Medical University, Taichung, Taiwan, R.O.C.;*

Abstract. Aim: To investigate the effect of oxidative stress biomarkers on the risk of potentially malignant oral disorders (PMODs). Materials and Methods: A total of 208 male adults with PMODs and an equal number of same-age control patients were enrolled. Plasma biomarkers of oxidative stress, measured with 8-hydroxy-2'-deoxyguanosine (8-OHdG) and 8-isoprostane (8-ISO), were determined using enzyme-linked immunosorbent assay kits. PMODs were diagnosed in accordance with the World Health Organization (WHO) guidelines. Results: A significant association between a high level of 8-ISO and an increased risk of PMODs was identified [odds ratio (OR)=1.71, 95% confidence interval (CI)=1.12-2.63; $p=0.013$]. This positive association was stronger among patients with PMOD subtype of leukoplakia (OR=1.94, 95% CI=1.24-3.06; $p=0.004$). However, no significant association was observed between plasma 8-OHdG levels and overall risk of PMODs or subtypes. Conclusion: Our findings suggest that increased plasma 8-ISO levels may indicate the prominence of lipid peroxidation in the development of PMODs, particularly leukoplakia.

Potentially malignant oral disorders (PMODs) pose a pressing public health challenge because of their high rate of transformation into oral malignancies. In Taiwan, oral cancer

(OC) is a leading cause of mortality, especially among men aged 24 to 44 years (1), indicating the necessity of early intervention.

Tumor development is a complex process that is mediated by numerous endogenous and exogenous factors (2). Prominent among these is the induction of reactive oxygen species (ROS). At normal physiological concentrations, ROS act as signal molecules mediating cell growth, migration, and differentiation. High concentrations, however, induce cell death, apoptosis, and senescence (3), particularly when the antioxidant defense is overwhelmed. ROS can oxidize lipids and proteins and interact directly with DNA, thereby triggering oxidative stress (4, 5), which can lead to manifestation of 8-hydroxy-2'-deoxyguanosine (8-OHdG) and 8-isoprostane (8-ISO) as biomarkers of oxidative damage (4, 6). Oxidative stress has been implicated in the pathogenesis of various pathologies, including oral tumors (7, 8).

The cytochrome *P-450* (CYP-450) and glutathione *S*-transferase (GST) families of enzymes, which are involved in the metabolism and detoxification of toxic endogenous and environmental agents (9), play key roles in combating toxicity and ROS generation through their metabolites (10). We demonstrated in a previous study that genetic polymorphism in *GSTM1* and *CYP1A1*2C* may increase susceptibility to PMODs, especially among smokers (11). Moreover, accumulated evidence has helped identify certain lifestyle factors, such as cigarette smoking, alcohol consumption, and betel-nut chewing, as risks for PMODs (12). Whether oxidative stress is associated with the risk of PMODs and whether such an association may be modified by genetic, environmental or lifestyle factors remains unclear.

Thus, the aim of this study was to investigate the association between oxidative stress biomarkers, measured based on 8-OHdG and 8-ISO levels, and the risk of PMODs.

Correspondences to: Professor Chih-Ching Yeh, Ph.D., MSPH, School of Public Health, College of Public Health, Taipei Medical University, 250 Wu-Hsing Street, Taipei 11031, Taiwan. Tel: +886 27361661 ext. 6534, Fax: +886 227384831, e-mail: ccyeh@tmu.edu.tw

Key Words: Potentially malignant oral disorders, leukoplakia, oxidative stress, biomarkers.

Table I. Comparison of characteristics of healthy controls and patients with potentially malignant oral disorders.

Variable ^a	Controls (N=208)	Cases (N=208)	p-Value ^c
Mean±SD age, years	38.45±9.3	38.78±9.1	0.719
Mean BMI±SD, kg/m ²	24.07±3.1	24.76±3.8	0.038
BMI			0.019
Normal weight (<24 kg/m ²)	116 (55.77%)	92 (44.23%)	
Overweight (≥24 kg/m ²)	92 (44.23%)	116 (55.77%)	
Educational level, n (%)			0.076
Below elementary school	35 (16.83%)	40 (19.51%)	
Junior high school	90 (43.27%)	105 (51.22%)	
Above senior high school	83 (39.90%)	60 (29.27%)	
Cigarette smoking, n (%)			0.141
Nonsmoker	11 (5.29%)	5 (2.40%)	
Current smoker	180 (86.54%)	192 (92.31%)	
Ex-smoker	17 (8.17%)	11 (5.29%)	
Cigarette pack-years, n (%) ^b			
<13	108 (51.92%)	96 (46.15%)	0.239
≥13	100 (48.08%)	112 (53.85%)	
Alcohol consumption, n (%)			0.871
Never	73 (35.10%)	68 (32.69%)	
Current drinker	75 (36.06%)	77 (37.02%)	
Ex-drinker	60 (28.85%)	63 (30.29%)	
Betel-nut chewing, n (%)			0.019
Never	76 (36.54%)	50 (24.04%)	
Current chewer	48 (23.08%)	62 (29.81%)	
Ex-chewer	84 (40.38%)	96 (46.15%)	
Family history of cancer, n (%)			0.817
No	160 (76.92%)	158 (75.96%)	
Yes	48 (23.08%)	50 (24.04%)	
Median 8-OHdG (range), ng/mg	51.59 (228)	43.33 (209)	0.053
Median 8-ISO (range), pg/ml	666.21 (3565)	790.26 (2242)	0.009

SD, Standard deviation; BMI, body-mass index; 8-OHdG, 8-hydroxy-2'-deoxyguanosine; 8-ISO, 8-isoprostane. ^aNumbers may not equal the total number because of missing data. ^bDivided according to median cigarette pack-years among control patients. ^cStudent's *t*-test, Pearson's Chi-squared test, or Kruskal–Wallis test.

Additionally, we investigated how lifestyle factors and genetic polymorphisms affected the association between oxidative stress biomarkers and PMOD risk.

Materials and Methods

Participants. This case–control study included 208 male patients with PMODs and an equal number of control patients without oral lesions. A detailed description of participant characteristics has already been published (11). In brief, participants were recruited from a male penitentiary in Taiwan. A complete oral examination was conducted on all participants by two experienced otolaryngologists in accordance with World Health Organization guidelines (13). Patients with PMOD were those diagnosed with submucous fibrosis or leukoplakia, and controls of the same ages were randomly selected from individuals with healthy oral-screening results.

All eligible subjects completed a self-administered questionnaire covering information on sociodemographics, personal and family medical history, and lifestyle habits such as cigarette smoking, betel-nut chewing, and alcohol consumption. Based on their habits,

participants were categorized as nonsmokers, current smokers (had been smoking for more than 1 year), or former smokers (had not smoked for at least the previous 6 months). Total cigarette consumption was computed in pack-years (PY), as the number of packs of cigarettes smoked daily multiplied by the duration of time (years) spent smoking. Body-mass index (BMI) was calculated from weight and height measurements as (weight/height²; kg/m²). The Ethics Review Committee of China Medical University and Hospital approved this study (IRB No: DMR96-IRB-085). All participants provided written, informed consent following a detailed explanation of the study objectives.

DNA isolation and genotyping. The methods for DNA isolation and genotyping were reported previously (11). In summary, 10 ml of venous blood was collected from all participants for genomic extraction. From buffy-coated cells, genomic DNA for genotyping was isolated. Genotyping of *GSTM1*, *GSTT1*, *GSTP1*, and *CYP1A1*2C* was performed through polymerase chain reaction/restriction fragment length polymorphism. A 10% random sample was retested and showed 100% concordance for all polymorphisms. All laboratory personnel were blinded to disease status of participants.

Table II. Associations between biomarkers of oxidative stress and risk of potentially malignant oral disorders.

Biomarker ^a	Controls		All cases		Leukoplakia			Submucous fibrosis		
	N=208	N=208	OR (95% CI) ^b	p-Value	N=168	OR (95% CI) ^b	p-Value	N=40	OR (95% CI) ^b	p-Value
8-OHdG, ng/mg										
<51.59	104	119	1.00		87	1.00		32	1.00	
≥51.59	104	89	0.71 (0.47-1.09)	0.115	81	0.93 (0.59-1.45)	0.744	8	0.22 (0.31-1.03)	0.063
8-ISO, pg/ml										
<666.21	104	79	1.00		59	1.00		20	1.00	
≥666.21	104	129	1.71 (1.12-2.63)	0.013	109	1.94 (1.24-3.06)	0.004	20	1.24 (0.58-2.67)	0.587

8-ISO, 8-Isoprostane; BMI, body-mass index; 8-OHdG, 8-hydroxy-2'-deoxyguanosine; OR, odds ratio; CI, confidence interval. ^aMedian values based on controls. ^bAdjusted for education, BMI, cigarette smoking (pack-years), alcohol consumption, and betel-nut chewing.

Measurements of plasma 8-OHdG and 8-ISO. Measurements of 8-OHdG and 8-ISO in plasma samples were carried out using enzyme-linked immunosorbent assay kits (Cayman Chemical Company, Ann Arbor, MI, USA) by following the kit protocol. The intra- and inter-assay variability coefficients were 4.7%-11.6% and 4.5%-10.7%, respectively, for 8-OHdG, and 6.4%-12.6% and 10.5%-24.3%, respectively, for 8-ISO.

Statistical analysis. Data analysis was performed using the SAS Statistical Package (Vers. 9.4 for Windows; SAS Institute, Cary, NC, USA). Statistical differences between cases and controls were assessed using the Student's *t*-test, Pearson's chi-squared test, or the Kruskal-Wallis test, where appropriate. Logistic regression models were used to calculate odds ratios (ORs) and 95% confidence intervals (CIs) to assess the association between oxidative stress biomarkers and risk of PMODs. Adjusted ORs were calculated using multivariate logistic regression models with adjustments for education, BMI, cigarette smoking (PY), alcohol consumption, and betel-nut chewing. Further analysis was conducted to examine the effects of lifestyle factor-biomarker and gene-biomarker interactions on the risk of PMODs. The levels of 8-OHdG and 8-ISO were dichotomized into binary variables according to median values for the controls. The joint effects of lifestyle factors (cigarette smoking, alcohol consumption, and betel-nut chewing), and genetic polymorphisms (*GSTM1*, *GSTT1*, *GSTP1*, and *CYP1A1*2C*) with biomarkers were also estimated using multivariate regression models with adjustments for covariates. Due to the high proportion of smokers in our sample, the median smoking time among controls (13 PY) was used to examine the interaction between biomarkers and smoking habit. Statistical significance was two-sided and set at $p < 0.05$.

Results

Table I compares characteristics between cases and controls. The case group had higher mean BMI ($p=0.038$), a higher proportion of overweight subjects ($p=0.019$), a higher proportion of current and former betel-nut chewers ($p=0.019$), a lower median 8-OHdG level ($p=0.053$), and a higher median 8-ISO level ($p=0.009$) than did the control group.

No significant association was observed between 8-OHdG and overall risk of PMODs or leukoplakia in the multivariate

logistic regression models. However, high levels of 8-OHdG were marginally significantly associated with lower risk of submucous fibrosis ($p=0.063$). Additionally, 8-ISO was significantly associated with an increased risk of PMODs ($p=0.013$) and leukoplakia ($p=0.004$) but not with an increased risk of submucous fibrosis ($p=0.587$) (Table II). Therefore, we further examined the joint effects of 8-ISO and lifestyle or genetic factors on the risk of PMODs.

Table III lists the effects of 8-ISO on the risk of PMODs and leukoplakia, as stratified according to lifestyle factors. No statistically significant interaction between 8-ISO and lifestyle factors was observed; however, 8-ISO elevation conferred an approximately two-fold or more increased risk of PMODs and leukoplakia in participants with ≤13 PY of smoking ($p=0.018$ and $p=0.002$, respectively) and in participants who ever consumed alcohol ($p=0.046$ and $p=0.014$, respectively). Moreover, 8-ISO elevation more than doubled the risk of leukoplakia alone in participants who had never been betel-nut chewers ($p=0.027$) and significantly increased risk in those who had been betel-nut chewers ($p=0.046$).

No significant interaction was observed between genetic polymorphisms and the oxidative stress biomarker on risks for PMODs or leukoplakia. After stratifying according to genotype, participants with *GSTM1*-null ($p=0.038$) or *CYP1A1*2C* AA ($p=0.034$) genotypes and high 8-ISO level exhibited a significantly increased risk of leukoplakia. Participants with *GSTT1*-null ($p=0.005$ and $p=0.002$) or *GSTP1* AA ($p=0.027$ and $p=0.015$) genotypes and high 8-ISO level exhibited increased risk of both PMOD and leukoplakia (Table IV).

Discussion

PMODs carry a significant risk for malignant transformation (14). Among them, OC is highly prevalent in Taiwan (15). Evidence has linked oxidative DNA damage and lipid peroxidation to such malignancies, including OC (7, 16, 17). In this study, we demonstrated that a high 8-ISO level was

Table III. Association between 8-isoprostane (8-ISO) level and the risk of potentially malignant oral disorders, stratified according to lifestyle factors.

		Controls N=208	All cases			Leukoplakia		
Factor	8-ISO, pg/ml		N=208	OR (95% CI)	p-Value	N=168	OR (95% CI)	p-Value
Cigarette smoking (pack-years)								
≤13	<666.21	56	34	1.00		21	1.00	
	≥666.21	52	62	2.07 (1.13-3.79) ^a	0.018	53	2.83 (1.45-5.53) ^a	0.002
>13	<666.21	48	45	1.00		38	1.00	
	≥666.21	52	67	1.47 (0.77-2.81) ^a	0.339	56	1.43 (0.74-2.78) ^a	0.288
p for interaction					0.408			0.099
Alcohol consumption								
Never	<666.21	35	24	1.00		21	1.00	
	≥666.21	38	44	2.05 (0.89-4.70) ^b	0.091	37	1.90 (0.81-4.47) ^b	0.141
Ever	<666.21	69	55	1.00		38	1.00	
	≥666.21	66	85	1.70 (1.01-2.87) ^b	0.046	72	2.03 (1.15-3.55) ^b	0.014
p for interaction					0.640			0.937
Betel-nut chewing								
Never	<666.21	38	20	1.00		15	1.00	
	≥666.21	38	30	2.12 (0.95-4.73) ^c	0.067	28	2.68 (1.12-6.43) ^c	0.027
Ever	<666.21	66	59	1.00		44	1.00	
	≥666.21	66	99	1.60 (0.95-2.69) ^c	0.077	81	1.75 (1.01-3.04) ^c	0.046
p for interaction					0.983			0.721

OR, Odds ratio; CI, confidence interval. ^aAdjusted for education, BMI, alcohol consumption, and betel-nut chewing. ^bAdjusted for education, BMI, cigarette smoking (pack-years), and betel-nut chewing. ^cAdjusted for education, BMI, alcohol consumption, and cigarette smoking (pack-years).

significantly associated with an increased risk of PMODs, particularly leukoplakia. We also revealed that both genetics and lifestyle factors may modify the effects of these biomarkers on the risk of PMODs.

As a byproduct of oxidative DNA damage, 8-OHdG is considered a reliable biomarker of oxidative damage (18). Research evidence has established a correlation between 8-OHdG level and PMODs and OC (8). Conversely, our analysis revealed lower levels of 8-OHdG in patients with PMODs than in controls, indicating that a high 8-OHdG level conferred no risk of PMODs. In this study, we obtained measurements of plasma 8-OHdG levels, that may reflect both oxidative DNA damage and a deficient DNA-repair system. A defect in the repair of 8-OHdG residue may indicate that 8-OHdG levels are elevated in tissue and peripheral leukocytes, but low in plasma (19). Thus, our results may reflect a defective DNA-repair system in our case patients because plasma 8-OHdG is derived from residual repaired 8-OHdG in DNA. The 8-OHdG levels measured in plasma were also lower in patients with colon cancer than in control patients (20). Similarly, Rossner *et al.* determined there was no association between plasma 8-OHdG level and breast cancer (21).

ROS-induced lipid peroxidation has been shown to be mutagenic and may serve as a reliable biomarker of both cellular damage and DNA adducts (8). Therefore, we also

assessed oxidative stress by examining plasma 8-ISO levels. Our data revealed that patients with high 8-ISO levels had increased susceptibility to PMODs and leukoplakia, which is consistent with the findings of other studies on patients with colon cancer, oesophageal cancer, and leukoplakia (17, 22, 23). The sensitivity of cell membranes to ROS induces lipid peroxidation that can produce a variety of mutagenic carbonyl products (24) and contribute to neoplastic transformation.

We further examined the combined effects of 8-ISO and lifestyle factors on the risk of PMODs and leukoplakia. No significant effects were observed, but individuals with a high 8-ISO level who fell in the category of either smokers of ≤13 PY or those who had ever consumed alcohol exhibited a significantly increased risk of both PMODs and leukoplakia. Alcohol consumption and smoking are established risk factors for PMODs (12). Interestingly, only those with high 8-ISO levels who had smoked for relatively few PY (≤13) exhibited an increased risk of PMODs and leukoplakia. This may be because the majority of our participants (96%) were cigarette smokers. When we stratified the results according to smoking status (never-smoked and ever-smoked), the results were significant for those who ever-smoked, but no results were generated for participants who had never smoked because of their small number. Since, we could not elucidate the effects of 8-ISO on PMODs in nonsmokers, further studies may be required. We observed that regardless

Table IV. Association between 8-isoprostane (8-ISO) levels and risk of potentially malignant oral disorders, stratified according to genetic polymorphism.

Polymorphism	8-ISO, pg/ml	Controls (N=208) Number	All cases (N=208)			Leukoplakia (N=168)		
			Number	OR (95% CI) ^a	<i>p</i> -Value	Number	OR (95% CI) ^a	<i>p</i> -Value
<i>GSTM1</i>								
Present	<666.21	51	32	1.00		24	1.00	
	≥666.21	51	57	1.72 (0.88-3.36)	0.115	47	1.92 (0.92-3.98)	0.081
Null	<666.21	53	47	1.00		35	1.00	
	≥666.21	53	72	1.66 (0.9–2.96)	0.087	62	1.91 (1.04-3.51)	0.038
	<i>p</i> for interaction				0.681			0.833
<i>GSTT1</i>								
Present	<666.21	45	36	1.00		26	1.00	
	≥666.21	44	42	0.96 (0.48-1.94)	0.918	34	1.18 (0.5-2.50)	0.663
Null	<666.21	59	43	1.00		33	1.00	
	≥666.21	60	87	2.31 (1.29-4.13)	0.005	75	2.63 (1.42-4.87)	0.002
	<i>p</i> for interaction				0.178			0.208
<i>GSTP1</i>								
AA	<666.21	71	52	1.00		39	1.00	
	≥666.21	67	88	1.82 (1.07-3.10)	0.027	72	2.02 (1.15-3.56)	0.015
AG/GG	<666.21	33	27	1.00		20	1.00	
	≥666.21	37	41	1.53 (0.67-3.48)	0.310	37	1.61 (0.67-3.89)	0.290
	<i>p</i> for interaction				0.315			0.394
<i>CYP1A1*2C</i>								
AA	<666.21	55	49	1.00		39	1.00	
	≥666.21	70	87	1.69 (0.97-2.93)	0.064	77	1.87 (1.05-3.34)	0.034
AG/GG	<666.21	49	30	1.00		20	1.00	
	≥666.21	34	42	1.66 (0.81-3.39)	0.165	32	1.83 (0.84-3.99)	0.131
	<i>p</i> for interaction				0.415			0.402

8-OHdG, 8-Hydroxy-2'-deoxyguanosine; OR, odds ratio; CI, confidence interval; *GSTM1*, glutathione S-transferase mu 1; *GSTT1*, glutathione S-transferase theta 1; *GSTP1*, glutathione S-transferase pi 1; *CYP1A1*2C*, cytochrome *P-450 1A1*2C*. ^aAdjusted for education, BMI, alcohol consumption, cigarette smoking (pack-years), and betel-nut chewing.

of betel-nut chewing habit, 8-ISO only increased the overall risk of PMODs, which supports the conclusions of other reports regarding the independent effects of both 8-ISO and betel-nut chewing on leukoplakia (12, 17).

Our data also demonstrated that polymorphisms of drug-metabolizing genes may modify the effects of 8-ISO on PMOD risk. Participants with high 8-ISO levels and *GSTT1*-null or *GSTP1* AA genotypes exhibited increased risk of both PMODs and leukoplakia, whereas *GSTM1*-null and *CYP1A1*2C* AA genotypes exhibited an increased susceptibility to leukoplakia. These findings support those of other studies that both GST and CYP independently affect the development of PMODs and leukoplakia (25, 26). The adverse effects of ROS through lipid peroxidation have previously been linked to OC and leukoplakia development (17, 26). The development of PMODs involves multiple processes, including molecular and cellular processes, which lead to tumor formation. These processes are mediated by ROS (27), which have been determined to initiate and promote tumor formation (8). In the presence of low antioxidant levels, ROS also attack the polyunsaturated fatty acids of fatty-acid membranes, causing lipid peroxidation that eventually

results in changes to cell-membrane structure and functionality. The polymorphisms of *GSTM1*, *GSTT1*, *GSTP1*, and *CYP1A1*2C* have been discovered to affect OC and PMODs (25). Various studies have also demonstrated the association of both *GSTM1*-null and *CYP1A1*2C* AA genotypes with increased risk of leukoplakia (11, 28).

Our study findings must be interpreted in light of several limitations. Firstly, we measured plasma 8-OHdG, which may reflect not only oxidative DNA stress but also DNA repair efficiency. Secondly, we did not include nutritional and psychological factors, which may have some effects on oral tumors. Thus, the exclusion of these confounding factors in future studies may be required in order to determine the effects of oxidative stress on PMODs. Finally, our participants were males from a penitentiary and therefore might not be representative of the general Taiwanese population.

Conclusion

Our findings revealed that an elevated level of plasma 8-ISO, but not 8-OHdG in patients with PMOD compared to control

patients might indicate the prominence of lipid peroxidation in the development of PMODs, particularly leukoplakia.

Acknowledgements

The Authors are grateful to study participants.

References

- Su C-C, Tsai K-Y, Hsu Y-Y, Lin Y-Y and Lian I-B: Chronic exposure to heavy metals and risk of oral cancer in Taiwanese males. *Oral Oncol* 46(8): 586-590, 2010.
- Swenberg JA, Lu K, Moeller BC, Gao L, Upton PB, Nakamura J and Starr TB: Endogenous versus exogenous DNA adducts: Their role in carcinogenesis, epidemiology, and risk assessment. *Toxicol Sci* 120(Suppl 1): S130-S145, 2011.
- Valko M, Rhodes CJ, Moncol J, Izakovic M and Mazur M: Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 160(1): 1-40, 2006.
- Marczynski B, Rozynek P, Kraus T, Schlosser S, Raithel HJ and Baur X: Levels of 8-hydroxy-2'-deoxyguanosine in DNA of white blood cells from workers highly exposed to asbestos in Germany. *Mutat Res* 468(2): 195-202, 2000.
- Xing F, Li Z, Sun A and Xing D: Reactive oxygen species promote chloroplast dysfunction and salicylic acid accumulation in fumonisin B1-induced cell death. *FEBS Lett* 587(14): 2164-2172, 2013.
- Thomson CA, Giuliano AR, Shaw JW, Rock CL, Ritenbaugh CK, Hakim IA, Hollenbach KA, Alberts DS and Pierce JP: Diet and biomarkers of oxidative damage in women previously treated for breast cancer. *Nutr Cancer* 51(2): 146-154, 2005.
- Ma N, Tagawa T, Hiraku Y, Murata M, Ding X and Kawanishi S: 8-Nitroguanine formation in oral leukoplakia, a premalignant lesion. *Nitric Oxide* 14(2): 137-143, 2006.
- Choudhari SK, Chaudhary M, Gadbail AR, Sharma A and Tekade S: Oxidative and antioxidative mechanisms in oral cancer and precancer: A review. *Oral Oncol* 50(1): 10-18, 2014.
- Sheweita SA: Drug-metabolizing enzymes: Mechanisms and functions. *Curr Drug Metab* 1(2): 107-132, 2000.
- Sosa V, Moliné T, Somoza R, Paciucci R, Kondoh H and Leonart ME: Oxidative stress and cancer: An overview. *Ageing Res Rev* 12(1): 376-390, 2013.
- Li YF, Sung FC, Tsai MH, Hua CH, Liu CS, Huang YT and Yeh CC: Interactions between cigarette smoking and polymorphisms of xenobiotic-metabolizing genes: The risk of oral leukoplakia. *Dis Markers* 34(4): 247-255, 2013.
- Kadashetti V, Shivakumar KM, Chaudhary M, Patil S, Gawande M and Hande A: Influence of risk factors on patients suffering from potentially malignant disorders and oral cancer: A case-control study. *J Oral Maxillofac Pathol* 21(3): 455-456, 2017.
- Warnakulasuriya S, Johnson NW and van der Waal I: Nomenclature and classification of potentially malignant disorders of the oral mucosa. *J Oral Pathol Med* 36(10): 575-580, 2007.
- Speight PM, Khurram SA and Kujan O: Oral potentially malignant disorders: Risk of progression to malignancy. *Oral Surg Oral Med Oral Pathol Oral Radiol* 125(6): 612-627, 2017.
- Chiang C-J, Lo W-C, Yang Y-W, You S-L, Chen C-J and Lai M-S: Incidence and survival of adult cancer patients in Taiwan, 2002-2012. *J Formos Med Assoc* 115(12): 1076-1088, 2016.
- Poulsen HE, Prieme H and Loft S: Role of oxidative DNA damage in cancer initiation and promotion. *Eur J Cancer Prev* 7(1): 9-16, 1998.
- Srivastava KC and Shrivastava D: Analysis of plasma lipid peroxidation and antioxidant enzymes status in patients of oral leukoplakia: A case-control study. *J Int Soc Prev Community Dent* 6(Suppl 3): S213-s218, 2016.
- Ziech D, Franco R, Georgakilas AG, Georgakila S, Malamou-Mitsi V, Schoneveld O, Pappa A and Panayiotidis MI: The role of reactive oxygen species and oxidative stress in environmental carcinogenesis and biomarker development. *Chem Biol Interact* 188(2): 334-339, 2010.
- Kondo S, Toyokuni S, Tanaka T, Hiai H, Onodera H, Kasai H and Imamura M: Overexpression of the *hOGG1* gene and high 8-hydroxy-2'-deoxyguanosine (8-OHdG) lyase activity in human colorectal carcinoma: Regulation mechanism of the 8-OHdG level in DNA. *Clin Cancer Res* 6(4): 1394-1400, 2000.
- Niedernhofer LJ, Daniels JS, Rouzer CA, Greene RE and Marnett LJ: Malondialdehyde, a product of lipid peroxidation, is mutagenic in human cells. *J Biol Chem* 278(33): 31426-31433, 2003.
- Rossner P Jr., Gammon MD, Terry MB, Agrawal M, Zhang FF, Teitelbaum SL, Eng SM, Gaudet MM, Neugut AI and Santella RM: Relationship between urinary 15-F2t-isoprostane and 8-oxodeoxyguanosine levels and breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 15(4): 639-644, 2006.
- Brys M, Morel A, Forma E, Krzeslak A, Wilkosz J, Rozanski W and Olas B: Relationship of urinary isoprostanes to prostate cancer occurrence. *Mol Cell Biochem* 372(1-2): 149-153, 2013.
- Freitas Bde J, Lloret GR, Visacri MB, Tuan BT, Amaral LS, Baldini D, de Sousa VM, de Castro LL, Aguiar JR, Pincinato Ede C, Mazzola PG and Moriel P: High 15-F2t-isoprostane levels in patients with a previous history of nonmelanoma skin cancer: The effects of supplementary antioxidant therapy. *Biomed Res Int* 2015: 963569, 2015.
- Benamira M, Johnson K, Chaudhary A, Bruner K, Tibbetts C and Marnett LJ: Induction of mutations by replication of malondialdehyde modified M13 DNA in *Escherichia coli*: Determination of the extent of DNA modification, genetic requirements for mutagenesis, and types of mutations induced. *Carcinogenesis* 16(1): 93-99, 1995.
- Liu H, Jia J, Mao X and Lin Z: Association of *CYP1A1* and *GSTM1* polymorphisms with oral cancer susceptibility: A meta-analysis. *Medicine* 94(27): e895, 2015.
- Chole RH, Patil RN, Basak A, Palandurkar K and Bhowate R: Estimation of serum malondialdehyde in oral cancer and precancer and its association with healthy individuals, gender, alcohol, and tobacco abuse. *J Cancer Res Ther* 6(4): 487-491, 2010.
- Valko M, Jomova K, Rhodes CJ, Kuca K and Musilek K: Redox- and non-redox-metal-induced formation of free radicals and their role in human disease. *Arch Toxicol* 90(1): 1-37, 2016.
- He P, Wei M, Wang Y and Liu Q: Associations among glutathione S-transferase T1, M1, and P1 polymorphisms and the risk of oral leukoplakia. *Genet Test Mol Biomarkers* 20(6): 312-321, 2016.

Received June 26, 2018

Revised July 11, 2018

Accepted July 13, 2018