Lowering and Stabilizing PSA Levels in Advanced-prostate Cancer Patients With Oral Methioninase

QINGHONG HAN and ROBERT M. HOFFMAN

AntiCancer, Inc., San Diego, CA, U.S.A.

Abstract. *Background/Aim:* Methionine addiction is a general and fundamental hallmark of cancer due to the excess use of methionine for transmethylation reactions, termed the “Hoffman Effect”. Methionine addiction has been shown to be a highly-effective target for cancer therapy by methionine restriction with oral recombinant methioninase (o-rMETase) in preclinical studies, including patient- derived orthotopic xenograft (PDOX) mouse models of cancer. A clinical study of o-rMETase as a supplement showed a 70% reduction of PSA levels in a patient with bone-metastatic prostate cancer. *Materials and Methods:* In the present study, two advanced prostate-cancer patients took o-rMETase as a supplement for approximately one month. *Results:* One of the patients taking o-rMETase showed a 38% reduction of PSA levels and the second patient showed a 20% PSA reduction. *Conclusion:* o-rMETase shows promise for treating patients with advanced prostate cancer.

Methionine addiction is a general and fundamental hallmark of cancer that greatly distinguishes all tested cancer types from normal cells, with respect to their high requirement levels for methionine (1-8). The methionine addiction of cancer cells is due to excess transmethylation reactions, compared to normal cells, which leads to methionine overuse and depletion of the cellular pools of free methionine and S-adenosylmethionine (7, 8), which is called the “Hoffman Effect” (9-11), a possibly stronger effect than the Warburg effect for glucose (12). The elevated transmethylation in cancer is due in part to the over-methylation of histone H-3 lysine marks (13, 14).

Targeting methionine addiction in cancer with recombinant methioninase (rMETase), cloned from *P. putida*

and expressed in *Escherichia coli* (15), has been shown to be effective against all major cancer types in mouse models of cancer, including patient-derived orthotopic xenograft (PDOX) mouse models (16-42).

Previous preliminary safety studies on *i.v.* PEGylated rMETase in macaque monkeys showed that rMETase depleted plasma methionine to <5 $\mu\text{mol/L}$, with transient anemia being the only side effect (43). PEGylation of rMETase was necessary to prevent anaphylaxis due to repeated treatment (43). *i.v.* injection of rMETase also caused rapid loss of the rMETase co-factor pyridoxal 5-phosphate (PLP) and thereby, a loss of its enzymatic activity (43).

A previous pilot Phase I clinical trial of o-rMETase was carried out to determine rMETase toxicity and the extent of methionine depletion in high-stage cancer patients. Circulating methionine levels were lowered to 0.1 μM by short-term *i.v.* rMETase treatment, without toxicity (44, 45).

The discovery that rMETase could be effectively administered orally (o-rMETase) (26), solved the problem of anaphylaxis and dissociation of PLP from rMETase, since o-rMETase exerts its effects in the gastro-intestinal tract without entering the circulation. o-rMETase has been shown to be highly effective against recalcitrant sarcoma, pancreatic cancer, colon cancer, and melanoma in PDOX mouse models (17-20, 24-26, 32, 33, 39).

Prostate cancer is the second-leading cause of death of North American men (46). Prostate-specific antigen (PSA) is a blood biomarker of prostate cancer and its progression (46). Androgen-deprivation therapy has been combined with docetaxel to improve survival of prostate cancer patients (46-49). However, survival of hormone-independent prostate cancer patients is low (47), and in need of improved therapy.

Due to the very low toxicity of o-rMETase, a pilot clinical study of o-rMETase, as a supplement, was initiated in 2020, with an advanced prostate-cancer patient with bone-metastatic disease. During a three-month period of receiving o-rMETase as a supplement, the patient had a 70% drop in his PSA levels, without any side effects (50). In the present study, we report a reduction in PSA levels in two additional advanced prostate cancer patients taking o-rMETase as a supplement, without side effects.

This article is freely accessible online.

Correspondence to: Robert M. Hoffman, Ph.D., AntiCancer, Inc. 7917 Ostrow St. San Diego, CA 92111, U.S.A. Tel: +1 8586542555. e-mail: all@anticancer.com

Key Words: Methioninase, oral, methionine addiction, prostate cancer, PSA, patients, response.

Materials and Methods

rMETase production and formulation. rMETase was fermented in recombinant *E. coli* and purified using column chromatography, as previously described (15). Pure methioninase was dissolved in phosphate-buffered saline (PBS) at 5 mg/ml (250 units), which comprised one dose for oral administration, to be taken after breakfast and dinner.

Methionine measurement. Methionine was measured in the plasma by high performance liquid chromatography as previously described (51).

Results

Patients. Two prostate-cancer patients were treated with o-rMETase twice a day with a dose of 250 units each time. Patient #1 was 55 years old, and had a 35 mm prostate cancer at diagnosis. The patient was initially treated with image-guided radiotherapy. This patient had a Gleason score of 4+3, with perineural invasion and extracapsular extension. Patient #2 was 90 years old, and was diagnosed with prostate cancer, with rapidly rising PSA levels.

PSA reduction by o-rMETase. Patient #1 had a 38.3% PSA drop within four weeks (Figure 1). Patient #2 had a 20.1% PSA drop within four weeks which then stabilized (Figure 2).

Circulating methionine reduction by o-rMETase. Patient #1 had a 42.7% drop in methionine levels during a 12-day period of taking o-rMETase (Figure 3).

Discussion

o-rMETase has been previously shown to cause a 70% drop in the PSA levels of a bone-metastatic prostate-cancer patient who was treated with o-rMETase as a supplement for 3 months, with a starting PSA value of over 2000 ng/ml (50).

Initially, rMETase was administered intra-venously or intra-peritoneally in mouse models (1, 21, 44, 45). However, primate studies showed that repeated *i.v.* infusion of rMETase caused anaphylaxis, which could be prevented by PEGylation of rMETase (43). In 2018, it was found that rMETase could be effectively administered orally (o-rMETase) (26), and subsequent studies showed that o-rMETase was effective in PDOX models against all major cancer types (17-20, 24-26, 32, 33, 39). o-rMETase does not appear to enter the blood stream, thereby eliminating the risk of anaphylaxis, as well as eliminating loss of activity due to dissociation of the rMETase co-factor PLP. o-rMETase functions in the gut to degrade methionine, which subsequently lowers the blood levels of methionine, since there is no longer a source of external methionine (26).

o-rMETase is therefore much safer than injecting rMETase into the blood stream (52), and has allowed the initial

Figure 1. PSA levels in patient #1 before and after receiving o-rMETase as a supplement for 21 days. The patient took two daily doses of 250 units of o-rMETase.

development of o-rMETase as a supplement for patients. The fact that the origin of o-rMETase was in *P. putida*, a soil bacterium that survives in environments with a wide range of pH, enabled o-rMETase to evolve into an acid-resistant enzyme that can survive the low pH of the stomach (1, 26).

The two prostate-cancer patients in the present report responded to o-rMETase as observed by the reduction in PSA levels (Figures 1 and 2). Patient #1 periodically fasted and was on a vegan diet, which may have affected his methionine and PSA levels.

The present dose of 250 units twice a day may be modified in the future in order to maximally lower the circulating methionine levels. Modifying the diet with low methionine-containing foods (53), may also be helpful along with altered dosing and scheduling of o-rMETase, to lower methionine and PSA levels even further. After optimization of the dose, schedule and diet, cancer patients can be treated with combinations of o-rMETase and chemotherapy, as methionine restriction and chemotherapy were found to be synergistic (6, 36). Most promising is combination chemotherapy along with o-rMETase for blockade of the methionine-methylation axis (54-56).

Conflicts of Interest

The Authors report no conflicts of interest in relation to this study.

Authors' Contributions

Qinghong Han produced the recombinant methioninase and analyzed the data. Robert M. Hoffman wrote the paper.

Acknowledgements

This paper is dedicated to the memory of John Mays, AR Moossa, MD, Sun Lee, MD, Professor Li Jiaxi and Masaki Kitajima, MD.

Figure 2. PSA levels in patient #2 before and during his treatment with o-rMETase, twice daily, 250 units/dose. Red arrow indicates the start of o-rMETase treatment.

Figure 3. Daily methionine levels in the plasma of patient #1 during treatment with o-rMETase, twice daily, 250 units/dose.

References

- 1 Hoffman RM: Development of recombinant methioninase to target the general cancer-specific metabolic defect of methionine dependence: A 40-year odyssey. *Expert Opin Biol Ther* 15(1): 21-31, 2015. PMID: 25439528. DOI: 10.1517/14712598.2015.963050
- 2 Hoffman RM and Erbe RW: High *in vivo* rates of methionine biosynthesis in transformed human and malignant rat cells auxotrophic for methionine. *Proc Natl Acad Sci USA* 73(5): 1523-1527, 1976. PMID: 179090. DOI: 10.1073/pnas.73.5.1523
- 3 Hoffman RM and Jacobsen SJ: Reversible growth arrest in simian virus 40-transformed human fibroblasts. *Proc Natl Acad Sci USA* 73(5): 1528-1532, 1976. PMID: 179091. DOI: 10.1073/pnas.73.5.1528

- Sci USA 77(12): 7306-7310, 1980. PMID: 6261250. DOI: 10.1073/pnas.77.12.7306
- 4 Guo H, Lishko VK, Herrera H, Groce A, Kubota T and Hoffman RM: Therapeutic tumor-specific cell cycle block induced by methionine starvation *in vivo*. Cancer Res 53(23): 5676-5679, 1993. PMID: 8242623.
- 5 Yano S, Li S, Han Q, Tan Y, Bouvet M, Fujiwara T and Hoffman RM: Selective methionine-induced trap of cancer cells in S/G2 phase visualized by FUCCI imaging confers chemosensitivity. Oncotarget 5(18): 8729-8736, 2014. PMID: 25238266. DOI: 10.18632/oncotarget.2369
- 6 Stern PH and Hoffman RM: Enhanced *in vitro* selective toxicity of chemotherapeutic agents for human cancer cells based on a metabolic defect. J Natl Cancer Inst 76(4): 629-639, 1986. PMID: 3457200. DOI: 10.1093/jnci/76.4.629
- 7 Stern PH and Hoffman RM: Elevated overall rates of transmethylation in cell lines from diverse human tumors. In Vitro 20(8): 663-670, 1984. PMID: 6500606. DOI: 10.1007/BF02619617
- 8 Coalson DW, Mecham JO, Stern PH and Hoffman RM: Reduced availability of endogenously synthesized methionine for S-adenosylmethionine formation in methionine-dependent cancer cells. Proc Natl Acad Sci USA 79(14): 4248-4251, 1982. PMID: 6289297. DOI: 10.1073/pnas.79.14.4248
- 9 Kaiser P: Methionine dependence of cancer. Biomolecules 10(4): 568, 2020. PMID: 32276408. DOI: 10.3390/biom10040568
- 10 Lauinger L and Kaiser P: Sensing and signaling of methionine metabolism. Metabolites 11(2): 83, 2021. PMID: 33572567. DOI: 10.3390/metabo11020083
- 11 Borrego SL, Fahrman J, Hou J, Lin DW, Tromberg BJ, Fiehn O and Kaiser P: Lipid remodeling in response to methionine stress in MDA-MBA-468 triple-negative breast cancer cells. J Lipid Res Feb 26:100056, 2021. Epub ahead of print. PMID: 33647277. DOI: 10.1016/j.jlr.2021.100056
- 12 Pirote B, Goldman S, Massager N, David P, Wikler D, Vandesteene A, Salmon I, Brothi J and Levivier M: Comparison of 18F-FDG and 11C-methionine for PET-guided stereotactic brain biopsy of gliomas. J Nucl Med 45(8): 1293-1298, 2004. PMID: 15299051.
- 13 Yamamoto J, Inubushi S, Han Q, Tashiro Y, Sun Y, Sugisawa N, Hamada K, Nishino H, Aoki Y, Miyake K, Matsuyama R, Bouvet M, Endo I and Hoffman RM: Cancer-specific overmethylation of histone H3 lysines is necessary for methionine addiction and malignancy. BioRxiv, 2020. DOI:10.1101/2020.12.04.412437
- 14 Yamamoto J, Han Q, Inubushi S, Sugisawa N, Hamada K, Nishino H, Miyake K, Kumamoto T, Matsuyama R, Bouvet M, Endo I and Hoffman RM: Histone methylation status of H3K4me3 and H3K9me3 under methionine restriction is unstable in methionine-addicted cancer cells, but stable in normal cells. Biochem Biophys Res Commun 533(4): 1034-1038, 2020. PMID: 33019978. DOI: 10.1016/j.bbrc.2020.09.108
- 15 Tan Y, Xu M, Tan X, Tan X, Wang X, Saikawa Y, Nagahama T, Sun X, Lenz M and Hoffman RM: Overexpression and large-scale production of recombinant L-methionine-alpha-deaminogamma-mercaptopmethane-lyase for novel anticancer therapy. Protein Expr Purif 9(2): 233-245, 1997. PMID: 9056489. DOI: 10.1006/prep.1996.0700
- 16 Igarashi K, Kawaguchi K, Kiyuna T, Miyake K, Murakami T, Yamamoto N, Hayashi K, Kimura H, Miwa S, Tsuchiya H and Hoffman RM: Effective metabolic targeting of human osteosarcoma cells *in vitro* and in orthotopic nude-mouse models with recombinant methioninase. Anticancer Res 37(9): 4807-4812, 2017. PMID: 28870899. DOI: 10.21873/anticancer.11887
- 17 Higuchi T, Han Q, Miyake K, Oshiro H, Sugisawa N, Tan Y, Yamamoto N, Hayashi K, Kimura H, Miwa S, Igarashi K, Bouvet M, Singh SR, Tsuchiya H and Hoffman RM: Combination of oral recombinant methioninase and decitabine arrests a chemotherapy-resistant undifferentiated soft-tissue sarcoma patient-derived orthotopic xenograft mouse model. Biochem Biophys Res Commun 523(1): 135-139, 2020. PMID: 31839218. DOI: 10.1016/j.bbrc.2019.12.024
- 18 Higuchi T, Sugisawa N, Yamamoto J, Oshiro H, Han Q, Yamamoto N, Hayashi K, Kimura H, Miwa S, Igarashi K, Tan Y, Kuchipudi S, Bouvet M, Singh SR, Tsuchiya H and Hoffman RM: The combination of oral-recombinant methioninase and azacitidine arrests a chemotherapy-resistant osteosarcoma patient-derived orthotopic xenograft mouse model. Cancer Chemother Pharmacol 85(2): 285-291, 2020. PMID: 31705268. DOI: 10.1007/s00280-019-03986-0
- 19 Oshiro H, Tome Y, Kiyuna T, Yoon SN, Lwin TM, Han Q, Tan Y, Miyake K, Higuchi T, Sugisawa N, Katsuya Y, Park JH, Zang Z, Razmjooei S, Bouvet M, Clary B, Singh SR, Kanaya F, Nishida K and Hoffman RM: Oral recombinant methioninase overcomes colorectal-cancer liver metastasis resistance to the combination of 5-fluorouracil and oxaliplatin in a patient-derived orthotopic xenograft mouse model. Anticancer Res 39(9): 4667-4671, 2019. PMID: 31519565. DOI: 10.21873/anticancer.13648
- 20 Higuchi T, Oshiro H, Miyake K, Sugisawa N, Han Q, Tan Y, Park J, Zhang Z, Razmjooei S, Yamamoto N, Hayashi K, Kimura H, Miwa S, Igarashi K, Bouvet M, Chawla SP, Singh SR, Tsuchiya H and Hoffman RM: Oral recombinant methioninase, combined with oral caffeine and injected cisplatin, overcome cisplatin-resistance and regresses patient-derived orthotopic xenograft model of osteosarcoma. Anticancer Res 39(9): 4653-4657, 2019. PMID: 31519563. DOI: 10.21873/anticancer.13646
- 21 Kawaguchi K, Han Q, Li S, Tan Y, Igarashi K, Murakami T, Unno M and Hoffman RM: Efficacy of recombinant methioninase (rMETase) on recalcitrant cancer patient-derived orthotopic xenograft (PDOX): A review. Cells 8(5): 410, 2019. PMID: 31052611. DOI: 10.3390/cells8050410
- 22 Miyake K, Kiyuna T, Li S, Han Q, Tan Y, Zhao M, Oshiro H, Kawaguchi K, Higuchi T, Zhang Z, Razmjooei S, Barangi M, Wangsiricharoen S, Murakami T, Singh AS, Li Y, Nelson SD, Eilber FC, Bouvet M, Hiroshima Y, Chishima T, Matsuyama R, Singh SR, Endo I and Hoffman RM: Combining tumor-selective bacterial therapy with Salmonella typhimurium A1-R and cancer metabolism targeting with oral recombinant methioninase regressed an Ewing's sarcoma in a patient-derived orthotopic xenograft model. Chemotherapy 63(5): 278-283, 2018. PMID: 30673664. DOI: 10.1159/000495574
- 23 Igarashi K, Kawaguchi K, Kiyuna T, Miyake K, Miyaki M, Yamamoto N, Hayashi K, Kimura H, Miwa S, Higuchi T, Singh AS, Chmielowski B, Nelson SD, Russell TA, Eckardt MA, Dry SM, Li Y, Singh SR, Chawla SP, Eilber FC, Tsuchiya H and Hoffman RM: Metabolic targeting with recombinant methioninase combined with palbociclib regresses a doxorubicin-resistant dedifferentiated liposarcoma. Biochem Biophys Res Commun 506(4): 912-917, 2018. PMID: 30392912. DOI: 10.1016/j.bbrc.2018.10.119

- 24 Higuchi T, Kawaguchi K, Miyake K, Han Q, Tan Y, Oshiro H, Sugisawa N, Zhang Z, Razmjooei S, Yamamoto N, Hayashi K, Kimura H, Miwa S, Igarashi K, Chawla SP, Singh AS, Eilber FC, Singh SR, Tsuchiya H and Hoffman RM: Oral recombinant methioninase combined with caffeine and doxorubicin induced regression of a doxorubicin-resistant synovial sarcoma in a PDOX mouse model. *Anticancer Res* 38(10): 5639-5644, 2018. PMID: 30275182. DOI: 10.21873/anticancer.12899
- 25 Kawaguchi K, Higuchi T, Li S, Han Q, Tan Y, Igarashi K, Zhao M, Miyake K, Kiyuna T, Miyake M, Ohshiro H, Sugisawa N, Zhang Z, Razmjooei S, Wangsiricharoen S, Chmielowski B, Nelson SD, Russell TA, Dry SM, Li Y, Eckardt MA, Singh AS, Singh SR, Eilber FC, Unno M and Hoffman RM: Combination therapy of tumor-targeting Salmonella typhimurium A1-R and oral recombinant methioninase regresses a BRAF-V600E-negative melanoma. *Biochem Biophys Res Commun* 503(4): 3086-3092, 2018. PMID: 30166061. DOI: 10.1016/j.bbrc.2018.08.097
- 26 Kawaguchi K, Miyake K, Han Q, Li S, Tan Y, Igarashi K, Kiyuna T, Miyake M, Higuchi T, Oshiro H, Zhang Z, Razmjooei S, Wangsiricharoen S, Bouvet M, Singh SR, Unno M and Hoffman RM: Oral recombinant methioninase (o-rMETase) is superior to injectable rMETase and overcomes acquired gemcitabine resistance in pancreatic cancer. *Cancer Lett* 432: 251-259, 2018. PMID: 29928962. DOI: 10.1016/j.canlet.2018.06.016
- 27 Kawaguchi K, Miyake K, Han Q, Li S, Tan Y, Igarashi K, Lwin TM, Higuchi T, Kiyuna T, Miyake M, Oshiro H, Bouvet M, Unno M and Hoffman RM: Targeting altered cancer methionine metabolism with recombinant methioninase (rMETase) overcomes partial gemcitabine-resistance and regresses a patient-derived orthotopic xenograft (PDOX) nude mouse model of pancreatic cancer. *Cell Cycle* 17(7): 868-873, 2018. PMID: 29623758. DOI: 10.1080/15384101.2018.1445907
- 28 Kawaguchi K, Han Q, Li S, Tan Y, Igarashi K, Miyake K, Kiyuna T, Miyake M, Chmielowski B, Nelson SD, Russell TA, Dry SM, Li Y, Singh AS, Eckardt MA, Unno M, Eilber FC and Hoffman RM: Intra-tumor L-methionine level highly correlates with tumor size in both pancreatic cancer and melanoma patient-derived orthotopic xenograft (PDOX) nude-mouse models. *Oncotarget* 9(13): 11119-11125, 2018. PMID: 29541401. DOI: 10.18632/oncotarget.24264
- 29 Kawaguchi K, Igarashi K, Li S, Han Q, Tan Y, Kiyuna T, Miyake K, Murakami T, Chmielowski B, Nelson SD, Russell TA, Dry SM, Li Y, Unno M, Eilber FC and Hoffman RM: Combination treatment with recombinant methioninase enables temozolomide to arrest a BRAF V600E melanoma in a patient-derived orthotopic xenograft (PDOX) mouse model. *Oncotarget* 8(49): 85516-85525, 2017. PMID: 29156737. DOI: 10.18632/oncotarget.20231
- 30 Igarashi K, Kawaguchi K, Kiyuna T, Miyake K, Miyake M, Li S, Han Q, Tan Y, Zhao M, Li Y, Nelson SD, Dry SM, Singh AS, Elliott IA, Russell TA, Eckardt MA, Yamamoto N, Hayashi K, Kimura H, Miwa S, Tsuchiya H, Eilber FC and Hoffman RM: Tumor-targeting Salmonella typhimurium A1-R combined with recombinant methioninase and cisplatin eradicates an osteosarcoma cisplatin-resistant lung metastasis in a patient-derived orthotopic xenograft (PDOX) mouse model: Decoy, trap and kill chemotherapy moves toward the clinic. *Cell Cycle* 17(6): 801-809, 2018. PMID: 29374999. DOI: 10.1080/15384101.2018.1431596
- 31 Igarashi K, Kawaguchi K, Li S, Han Q, Tan Y, Murakami T, Kiyuna T, Miyake K, Miyake M, Singh AS, Eckardt MA, Nelson SD, Russell TA, Dry SM, Li Y, Yamamoto N, Hayashi K, Kimura H, Miwa S, Tsuchiya H, Singh SR, Eilber FC and Hoffman RM: Recombinant methioninase in combination with doxorubicin (DOX) overcomes first-line DOX resistance in a patient-derived orthotopic xenograft nude-mouse model of undifferentiated spindle-cell sarcoma. *Cancer Lett* 417: 168-173, 2018. PMID: 29306021. DOI: 10.1016/j.canlet.2017.12.028
- 32 Kawaguchi K, Han Q, Li S, Tan Y, Igarashi K, Kiyuna T, Miyake K, Miyake M, Chmielowski B, Nelson SD, Russell TA, Dry SM, Li Y, Singh AS, Eckardt MA, Unno M, Eilber FC and Hoffman RM: Targeting methionine with oral recombinant methioninase (o-rMETase) arrests a patient-derived orthotopic xenograft (PDOX) model of BRAF-V600E mutant melanoma: implications for chronic clinical cancer therapy and prevention. *Cell Cycle* 17(3): 356-361, 2018. PMID: 29187018. DOI: 10.1080/15384101.2017.1405195
- 33 Park JH, Han Q, Zhao M, Tan Y, Higuchi T, Yoon SN, Sugisawa N, Yamamoto J, Bouvet M, Clary B, Singh SR and Hoffman RM: Oral recombinant methioninase combined with oxaliplatin and 5-fluorouracil regressed a colon cancer growing on the peritoneal surface in a patient-derived orthotopic xenograft mouse model. *Tissue Cell* 61: 109-114, 2019. PMID: 31759402. DOI: 10.1016/j.tice.2019.09.006
- 34 Igarashi K, Li S, Han Q, Tan Y, Kawaguchi K, Murakami T, Kiyuna T, Miyake K, Li Y, Nelson SD, Dry SM, Singh AS, Elliott IA, Russell TA, Eckardt MA, Yamamoto N, Hayashi K, Kimura H, Miwa S, Tsuchiya H, Eilber FC and Hoffman RM: Growth of doxorubicin-resistant undifferentiated spindle-cell sarcoma PDOX is arrested by metabolic targeting with recombinant methioninase. *J Cell Biochem* 119(4): 3537-3544, 2018. PMID: 29143983. DOI: 10.1002/jcb.26527
- 35 Murakami T, Li S, Han Q, Tan Y, Kiyuna T, Igarashi K, Kawaguchi K, Hwang HK, Miyake K, Singh AS, Nelson SD, Dry SM, Li Y, Hiroshima Y, Lwin TM, DeLong JC, Chishima T, Tanaka K, Bouvet M, Endo I, Eilber FC and Hoffman RM: Recombinant methioninase effectively targets a Ewing's sarcoma in a patient-derived orthotopic xenograft (PDOX) nude-mouse model. *Oncotarget* 8(22): 35630-35638, 2017. PMID: 28404944. DOI: 10.18632/oncotarget.15823
- 36 Yano S, Takehara K, Zhao M, Tan Y, Han Q, Li S, Bouvet M, Fujiwara T and Hoffman RM: Tumor-specific cell-cycle decoy by Salmonella typhimurium A1-R combined with tumor-selective cell-cycle trap by methioninase overcome tumor intrinsic chemoresistance as visualized by Fucci imaging. *Cell Cycle* 15(13): 1715-1723, 2016. PMID: 27152859. DOI: 10.1080/15384101.2016.1181240
- 37 Igarashi K, Kawaguchi K, Li S, Han Q, Tan Y, Gainor E, Kiyuna T, Miyake K, Miyake M, Higuchi T, Oshiro H, Singh AS, Eckardt MA, Nelson SD, Russell TA, Dry SM, Li Y, Yamamoto N, Hayashi K, Kimura H, Miwa S, Tsuchiya H, Eilber FC and Hoffman RM: Recombinant methioninase combined with doxorubicin (DOX) regresses a DOX-resistant synovial sarcoma in a patient-derived orthotopic xenograft (PDOX) mouse model. *Oncotarget* 9(27): 19263-19272, 2018. PMID: 29721200. DOI: 10.18632/oncotarget.24996
- 38 Kawaguchi K, Igarashi K, Li S, Han Q, Tan Y, Miyake K, Kiyuna T, Miyake M, Murakami T, Chmielowski B, Nelson SD, Russell TA, Dry SM, Li Y, Unno M, Eilber FC and Hoffman RM: Recombinant methioninase (rMETase) is an effective therapeutic for BRAF-V600E-negative as well as -positive melanoma in patient-derived orthotopic xenograft (PDOX) mouse models. *Oncotarget* 9(1): 915-923, 2017. PMID: 29416666. DOI: 10.18632/oncotarget.23185

- 39 Park JH, Zhao M, Han Q, Sun Y, Higuchi T, Sugisawa N, Yamamoto J, Singh SR, Clary B, Bouvet M and Hoffman RM: Efficacy of oral recombinant methioninase combined with oxaliplatin and 5-fluorouracil on primary colon cancer in a patient-derived orthotopic xenograft mouse model. *Biochem Biophys Res Commun* 518(2): 306-310, 2019. PMID: 31421825. DOI: 10.1016/j.bbrc.2019.08.051
- 40 Yoshioka T, Wada T, Uchida N, Maki H, Yoshida H, Ide N, Kasai H, Hojo K, Shono K, Maekawa R, Yagi S, Hoffman RM and Sugita K: Anticancer efficacy *in vivo* and *in vitro*, synergy with 5-fluorouracil, and safety of recombinant methioninase. *Cancer Res* 58(12): 2583-2587, 1998. PMID: 9635582.
- 41 Kokkinakis DM, Hoffman RM, Frenkel EP, Wick JB, Han Q, Xu M, Tan Y and Schold SC: Synergy between methionine stress and chemotherapy in the treatment of brain tumor xenografts in athymic mice. *Cancer Res* 61(10): 4017-4023, 2001. PMID: 11358820.
- 42 Tan Y, Sun X, Xu M, Tan X, Sasson A, Rashidi B, Han Q, Tan X, Wang X, An Z, Sun FX and Hoffman RM: Efficacy of recombinant methioninase in combination with cisplatin on human colon tumors in nude mice. *Clin Cancer Res* 5(8): 2157-2163, 1999. PMID: 10473100.
- 43 Yang Z, Wang J, Lu Q, Xu J, Kobayashi Y, Takakura T, Takimoto A, Yoshioka T, Lian C, Chen C, Zhang D, Zhang Y, Li S, Sun X, Tan Y, Yagi S, Frenkel EP and Hoffman RM: PEGylation confers greatly extended half-life and attenuated immunogenicity to recombinant methioninase in primates. *Cancer Res* 64(18): 6673-6678, 2004. PMID: 15374983. DOI: 10.1158/0008-5472.CAN-04-1822
- 44 Tan Y, Zavala J Sr, Han Q, Xu M, Sun X, Tan X, Tan X, Magana R, Geller J and Hoffman RM: Recombinant methioninase infusion reduces the biochemical endpoint of serum methionine with minimal toxicity in high-stage cancer patients. *Anticancer Res* 17(5B): 3857-3860, 1997. PMID: 9427792.
- 45 Tan Y, Zavala J Sr, Xu M, Zavala J Jr and Hoffman RM: Serum methionine depletion without side effects by methioninase in metastatic breast cancer patients. *Anticancer Res* 16(6C): 3937-3942, 1996. PMID: 9042316.
- 46 Sartor O and de Bono JS: Metastatic Prostate Cancer. *N Engl J Med* 378(7): 645-657, 2018. PMID: 29412780. DOI: 10.1056/NEJMr1701695
- 47 Siegel RL, Miller KD and Jemal A: Cancer statistics, 2020. *CA Cancer J Clin* 70(1): 7-30, 2020. PMID: 31912902. DOI: 10.3322/caac.21590
- 48 Sweeney CJ, Chen YH, Carducci M, Liu G, Jarrard DF, Eisenberger M, Wong YN, Hahn N, Kohli M, Cooney MM, Dreicer R, Vogelzang NJ, Picus J, Shevrin D, Hussain M, Garcia JA and DiPaola RS: Chemohormonal therapy in metastatic hormone-sensitive prostate cancer. *N Engl J Med* 373(8): 737-746, 2015. PMID: 26244877. DOI: 10.1056/NEJMoa1503747
- 49 James ND, Sydes MR, Clarke NW, Mason MD, Dearnaley DP, Spears MR, Ritchie AW, Parker CC, Russell JM, Attard G, de Bono J, Cross W, Jones RJ, Thalmann G, Amos C, Matheson D, Millman R, Alzouebi M, Beesley S, Birtle AJ, Brock S, Cathomas R, Chakraborti P, Chowdhury S, Cook A, Elliott T, Gale J, Gibbs S, Graham JD, Hetherington J, Hughes R, Laing R, McKinna F, McLaren DB, O'Sullivan JM, Parikh O, Peedell C, Protheroe A, Robinson AJ, Srihari N, Srinivasan R, Staffurth J, Sundar S, Tolan S, Tsang D, Wagstaff J, Parmar MK and STAMPEDE investigators: Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): Survival results from an adaptive, multiarm, multistage, platform randomised controlled trial. *Lancet* 387(10024): 1163-1177, 2016. PMID: 26719232. DOI: 10.1016/S0140-6736(15)01037-5
- 50 Han Q, Tan Y and Hoffman RM: Oral dosing of recombinant methioninase is associated with a 70% drop in PSA in a patient with bone-metastatic prostate cancer and 50% reduction in circulating methionine in a high-stage ovarian cancer patient. *Anticancer Res* 40(5): 2813-2819, 2020. PMID: 32366428. DOI: 10.21873/anticancer.14254
- 51 Sun X, Tan Y, Yang Z, Li S and Hoffman RM: A rapid HPLC method for the measurement of ultra-low plasma methionine concentrations applicable to methionine depletion therapy. *Anticancer Res* 25(1A): 59-62, 2005. PMID: 15816519.
- 52 Lu WC, Saha A, Yan W, Garrison K, Lamb C, Pandey R, Irani S, Lodi A, Lu X, Tiziani S, Zhang YJ, Georgiou G, DiGiovanni J and Stone E: Enzyme-mediated depletion of serum l-Met abrogates prostate cancer growth *via* multiple mechanisms without evidence of systemic toxicity. *Proc Natl Acad Sci USA* 117(23): 13000-13011, 2020. PMID: 32434918. DOI: 10.1073/pnas.1917362117
- 53 Gao X, Sanderson SM, Dai Z, Reid MA, Cooper DE, Lu M, Richie JP Jr, Ciccarella A, Calcagnotto A, Mikhael PG, Mentch SJ, Liu J, Ables G, Kirsch DG, Hsu DS, Nichenametla SN and Locasale JW: Dietary methionine influences therapy in mouse cancer models and alters human metabolism. *Nature* 572(7769): 397-401, 2019. PMID: 31367041. DOI: 10.1038/s41586-019-1437-3
- 54 Sugisawa N, Yamamoto J, Han Q, Tan Y, Tashiro Y, Nishino H, Inubushi S, Hamada K, Kawaguchi K, Unno M, Bouvet M and Hoffman RM: Triple-methyl blockade with recombinant methioninase, cycloleucine, and azacitidine arrests a pancreatic cancer patient-derived orthotopic xenograft model. *Pancreas* 50(1): 93-98, 2021. PMID: 33370029. DOI: 10.1097/MPA.0000000000001709
- 55 Higuchi T, Han Q, Sugisawa N, Yamamoto J, Yamamoto N, Hayashi K, Kimura H, Miwa S, Igarashi K, Bouvet M, Singh SR, Tsuchiya H and Hoffman RM: Combination methionine-methylation-axis blockade: A novel approach to target the methionine addiction of cancer. *Cancer Genomics Proteomics* 18(2): 113-120, 2021. PMID: 33608308. DOI: 10.21873/cgp.20246
- 56 Kalev P, Hyer ML, Gross S, Konteatis Z, Chen CC, Fletcher M, Lein M, Aguado-Fraile E, Frank V, Barnett A, Mandley E, Goldford J, Chen Y, Sellers K, Hayes S, Lizotte K, Quang P, Tuncay Y, Clasquin M, Peters R, Weier J, Simone E, Murtie J, Liu W, Nagaraja R, Dang L, Sui Z, Biller SA, Travins J, Marks KM and Marjon K: MAT2A inhibition blocks the growth of MTAP-deleted cancer cells by reducing PRMT5-dependent mRNA splicing and inducing DNA damage. *Cancer Cell* 39: 209-224, 2021. PMID: 33450196. DOI: 10.1016/j.ccell.2020.12.010

Received February 22, 2021

Revised March 8, 2021

Accepted March 11, 2021